

5-2016

Finite Involution Semigroups with Infinite Irredundant Bases of Identities

Edmond W. H. Lee

Nova Southeastern University, edmond.lee@nova.edu

Follow this and additional works at: https://nsuworks.nova.edu/math_facarticles



Part of the [Mathematics Commons](#)

NSUWorks Citation

Lee, Edmond W. H., "Finite Involution Semigroups with Infinite Irredundant Bases of Identities" (2016). *Mathematics Faculty Articles*. 150.

https://nsuworks.nova.edu/math_facarticles/150

This Article is brought to you for free and open access by the Department of Mathematics at NSUWorks. It has been accepted for inclusion in Mathematics Faculty Articles by an authorized administrator of NSUWorks. For more information, please contact nsuworks@nova.edu.

Research Article

Edmond W. H. Lee

Finite involution semigroups with infinite irredundant bases of identities

Abstract: A basis of identities for an algebra is *irredundant* if each of its proper subsets fails to be a basis for the algebra. The first known examples of finite involution semigroups with infinite irredundant bases are exhibited. These involution semigroups satisfy several counterintuitive properties: their semigroup reducts do not have irredundant bases, they share reducts with some other finitely based involution semigroups, and they are direct products of finitely based involution semigroups.

Keywords: Semigroup, involution semigroup, identity, basis, irredundant basis, finite basis

MSC 2010: 20M05

Edmond W. H. Lee: Division of Math, Science, and Technology, Nova Southeastern University, 3301 College Avenue, Fort Lauderdale, Florida 33314, USA, e-mail: edmond.lee@nova.edu

Communicated by: Manfred Droste

1 Introduction

1.1 Finite basis problem

A *basis* for an algebra A is a set of identities satisfied by A that axiomatizes all identities of A . An algebra is *finitely based* if it has some finite basis. Finite groups [16], finite associative rings [7, 12], and finite Lie algebras [1] are all finitely based. However, this result does not hold for all finite algebras; there exist groupoids with as few as three elements that are not finitely based [5, 15]. In general, the finite basis problem for finite algebras is undecidable [14], but the problem remains open when restricted to finite semigroups. The first examples of non-finitely based finite semigroups [17] were discovered in the 1960s. Since then, the finite basis problem for finite semigroups has been intensely investigated. For further information, refer to the survey by Volkov [19].

1.2 Irredundant bases

A basis for an algebra A is *irredundant* if each of its proper subsets fails to be a basis for A . An algebra is *irredundantly based* if it has some irredundant basis. It is clear that any finitely based algebra is irredundantly based. As for non-finitely based finite algebras, there was initial hope that they all have irredundant bases, but this was refuted by subsequent examples of non-irredundantly based finite semigroups [13, 18]. On the other hand, there also exist finite semigroups with infinite irredundant bases [4]. Apart from these examples, not much is known about the irredundant basis property in general. The problem of deciding if a non-finitely based finite algebra is irredundantly based remains open.

1.3 Involution semigroups

Recall that an *involution semigroup* is a pair $(S, *)$, where the *reduct* S is a semigroup and $x \mapsto x^*$ is a unary operation on S such that

$$(x^*)^* = x \quad \text{and} \quad (xy)^* = y^*x^*.$$

Examples of involution semigroups include any group with inversion $x \mapsto x^{-1}$ and the multiplicative semigroup of all $n \times n$ matrices over any field with transposition $x \mapsto x^T$. Compared with semigroups, even less is known about involution semigroups with respect to the irredundant basis property. Presently, only one finite involution semigroup is known to be non-irredundantly based [6]. Explicit examples of involution semigroups with infinite irredundant bases are unavailable. The objective of the present article is to exhibit the first examples of such finite involution semigroups. These new examples, together with recently established results [9, 10], demonstrate that an involution semigroup and its reduct can satisfy very contrasting equational properties.

1.4 Known results

The main examples of the present article are constructed from the cyclic group

$$\mathbb{Z}_n = \langle g \mid g^n = 1 \rangle = \{1, g, g^2, \dots, g^{n-1}\}$$

of order $n \geq 1$ and the \mathcal{J} -trivial semigroup

$$L = \langle e, f \mid e^2 = e, f^2 = f, efe = 0 \rangle = \{0, e, f, ef, fe, fef\}$$

of order six. The multiplication table of L is given as follows.

L	0	fef	ef	fe	e	f
0	0	0	0	0	0	0
fef	0	0	0	0	0	fef
ef	0	0	0	0	0	ef
fe	0	0	fef	0	fe	fef
e	0	0	ef	0	e	ef
f	0	fef	fef	fe	fe	f

Up to isomorphism, the semigroup L is one of only four non-finitely based semigroups of order six [11]. The direct product $L \times \mathbb{Z}_n$ of L and \mathbb{Z}_n is a non-finitely based semigroup [8], and this result was recently generalized.

Theorem 1.1 ([9, Theorem 1.3]). *For each $n \geq 1$, the non-finitely based semigroup $L \times \mathbb{Z}_n$ is non-irredundantly based.*

For each $n \geq 1$, let $\text{sq}(n)$ denote the set of all square roots of unity modulo n , that is,

$$\text{sq}(n) = \{\mathbb{R} \in \{1, 2, \dots, n\} : \mathbb{R}^2 \equiv 1 \pmod{n}\}.$$

It is routinely checked that for each $\mathbb{R} \in \text{sq}(n)$,

$$\mathbb{Z}_n^{\mathbb{R}} = (\mathbb{Z}_n, \mathbb{R}) \quad \text{with} \quad x \mapsto x^{\mathbb{R}}$$

is an involution semigroup. Conversely, any involution semigroup with reduct \mathbb{Z}_n is isomorphic to some $\mathbb{Z}_n^{\mathbb{R}}$, so the number of such involution semigroups is

$$|\text{sq}(n)| = \begin{cases} 2^{\omega(n)+1} & \text{if } n \equiv 0 \pmod{8}, \\ 2^{\omega(n)-1} & \text{if } n \equiv \pm 2 \pmod{8}, \\ 2^{\omega(n)} & \text{otherwise,} \end{cases}$$

where $\omega(n)$ is the number of distinct prime factors of n ([3]). In particular, the inversion operation $x \mapsto x^{-1}$ in \mathbb{Z}_n coincides with $x \mapsto x^{n-1}$.

As for the semigroup L , it is the reduct of the involution semigroup

$$\mathbf{L} = (L, \lambda),$$

where λ interchanges ef and fe but fixes all other elements of L . In fact, up to isomorphism, L is the unique involution semigroup with reduct L . Hence, the direct product of the involution semigroups L and $Z_n^{\mathbb{R}}$ is

$$L \times Z_n^{\mathbb{R}} = (L \times Z_n, *) \quad \text{with} \quad (x, y)^* = (x^\lambda, y^{\mathbb{R}}).$$

Consequently, up to isomorphism, the number of involution semigroups with reduct $L \times Z_n$ is precisely $|\text{sq}(n)|$.

The finite basis property of the involution semigroup $L \times Z_n^{\mathbb{R}}$, when $\mathbb{R} = 1$, was recently established.

Theorem 1.2 ([10, Theorem 1.2]). *For each $n \geq 1$, the involution semigroup $L \times Z_n^1$ is finitely based and so is irredundantly based.*

1.5 Main result

In view of Theorem 1.2, what remains to be investigated is the involution semigroup $L \times Z_n^{\mathbb{R}}$ with $\mathbb{R} \in \text{sq}(n) \setminus \{1\}$.

Theorem 1.3. *For each $n \geq 1$ and $\mathbb{R} \in \text{sq}(n) \setminus \{1\}$, the involution semigroup $L \times Z_n^{\mathbb{R}}$ has an infinite irredundant basis and so is non-finitely based.*

The results of Theorems 1.1–1.3 are summarized in Table 1.

	$L \times Z_n$	$L \times Z_n^1$	$L \times Z_n^{\mathbb{R}}, \mathbb{R} \in \text{sq}(n) \setminus \{1\}$
Finitely based	No	Yes	No
Irredundantly based	No	Yes	Yes

Table 1. Summary of Theorems 1.1–1.3.

Corollary 1.4. *For each $n \geq 1$ and $\mathbb{R} \in \text{sq}(n)$, the involution semigroup $L \times Z_n^{\mathbb{R}}$ is*

- (i) *irredundantly based,*
- (ii) *finitely based if and only if $\mathbb{R} = 1$.*

Corollary 1.5. *There exist finite examples of*

- (i) *involution semigroups with infinite irredundant bases such that their reducts do not have such bases,*
- (ii) *non-finitely based involution semigroups sharing the same reducts with finitely based involution semigroups,*
- (iii) *non-finitely based involution semigroups that are direct products of finitely based involution semigroups.*

Proof. Refer to Table 1 for examples of parts (i) and (ii). As for part (iii), the involution semigroup $L \times Z_n^{n-1}$ is non-finitely based for all $n \geq 3$, while the involution semigroups L and Z_n^{n-1} are finitely based. □

Since all finitely based algebras are irredundantly based, no involution semigroup can provide a $\begin{pmatrix} \text{Yes} \\ \text{No} \end{pmatrix}$ -column in Table 1. Therefore, this table exhibits the most complicated scenario for a non-irredundantly based finite semigroup S in the following sense: there exist involution semigroups $(S, *)$ and (S, \dagger) with reduct S such that no two of S , $(S, *)$, and (S, \dagger) simultaneously satisfy the same finite basis property and irredundant basis property. It is of interest to know if the most complicated scenario exists for an irredundantly based finite semigroup.

Question 1.6. *Is there an irredundantly based finite semigroup S , being the reduct of involution semigroups $(S, *)$ and (S, \dagger) , such that their finite basis property and irredundant basis property are described in one of the following tables?*

	S	$(S, *)$	(S, \dagger)
Finitely based	Yes	No	No
Irredundantly based	Yes	Yes	No

	S	$(S, *)$	(S, \dagger)
Finitely based	No	Yes	No
Irredundantly based	Yes	Yes	No

1.6 Smaller examples

Although the order $6n$ of the irredundantly based involution semigroup $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$ in Theorem 1.3 can be considered small, smaller examples are available. For each $n \geq 1$, the subset

$$L_n = (L \times \{1\}) \cup (\{0\} \times \mathbb{Z}_n)$$

of $L \times \mathbb{Z}_n$ has $n + 5$ elements. It is routinely verified that L_n is a subsemigroup of $L \times \mathbb{Z}_n$ that is closed under the unary operation $(x, y)^* \mapsto (x^\lambda, y^{\mathbb{R}})$ of $L \times \mathbf{Z}_n^{\mathbb{R}}$. Hence, $\mathbf{L}_n^{\mathbb{R}} = (L_n, *)$ is an involution subsemigroup of $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$ of order $n + 5$.

Lemma 1.7. *For each $n \geq 1$, the involution semigroups $\mathbf{L}_n^{\mathbb{R}}$ and $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$ satisfy the same identities.*

Proof. The identities satisfied by $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$ are vacuously satisfied by its involution subsemigroup $\mathbf{L}_n^{\mathbb{R}}$. Conversely, since

$$\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}} = (L, \lambda) \times (\mathbb{Z}_n, \mathbb{R}) \cong (L \times \{1\}, *) \times (\{0\} \times \mathbb{Z}_n, *) \subseteq \mathbf{L}_n^{\mathbb{R}} \times \mathbf{L}_n^{\mathbb{R}},$$

the identities satisfied by $\mathbf{L}_n^{\mathbb{R}}$ are also satisfied by $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$. □

Consequently, the involution semigroup \mathbf{L}_3^2 of order eight has an infinite irredundant basis; it is presently the smallest known example. As for semigroups with infinite irredundant bases, the smallest example currently known is of order nine [4].

Question 1.8. What is the smallest possible order of a semigroup or involution semigroup with infinite irredundant basis?

1.7 Organization

There are seven sections in the present article. Notation and background material are given in Section 2. Sections 3–6 are devoted to establishing an explicit infinite basis for the involution semigroup $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$. This explicit basis is then shown in Section 7 to contain an infinite irredundant basis for $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$. Consequently, the proof of Theorem 1.3 is complete.

2 Preliminaries

The *free involution semigroup* over a countably infinite alphabet \mathcal{A} is the free semigroup $(\mathcal{A} \cup \mathcal{A}^*)^+$ over the disjoint union of \mathcal{A} and $\mathcal{A}^* = \{x^* : x \in \mathcal{A}\}$, with endowed unary operation $*$ defined by $(x^*)^* = x$ for all $x \in \mathcal{A}$ and

$$(x_1 x_2 \cdots x_m)^* = x_m^* x_{m-1}^* \cdots x_1^*$$

for all $x_1, x_2, \dots, x_m \in \mathcal{A} \cup \mathcal{A}^*$. Elements of $\mathcal{A} \cup \mathcal{A}^*$ are called *letters* while elements of $(\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ are called *words*. Words in $\mathcal{A}^+ \cup \{\emptyset\}$ are called *plain words*. A word that is also a single letter is called a *singleton*. Any word $w \in (\mathcal{A} \cup \mathcal{A}^*)^+$ can be written in the form

$$w = x_1^{\otimes_1} x_2^{\otimes_2} \cdots x_m^{\otimes_m},$$

where $x_1, x_2, \dots, x_m \in \mathcal{A}$ and $\otimes_1, \otimes_2, \dots, \otimes_m \in \{1, *\}$; the *plain projection* of such a word is the plain word

$$\bar{w} = x_1 x_2 \cdots x_m.$$

The first and last letters that occur in a word w are denoted by $\triangleright w$ and $w \triangleleft$, respectively. The number of times that a letter x occurs in a word w is denoted by $\text{occ}(x, w)$. The *content* of a word w is defined to be the set

$$\text{con}(\bar{w}) = \{x \in \mathcal{A} : \text{occ}(x, \bar{w}) \geq 1\};$$

this set is partitioned into the sets

$$\text{sim}(\bar{w}) = \{x \in \mathcal{A} : \text{occ}(x, \bar{w}) = 1\} \quad \text{and} \quad \text{non}(\bar{w}) = \{x \in \mathcal{A} : \text{occ}(x, \bar{w}) \geq 2\}.$$

The letters in $\text{sim}(\bar{w})$ are said to be *simple* in w while the letters in $\text{non}(\bar{w})$ are said to be *non-simple* in w . A word w is *non-simple* if some letter is non-simple in it, that is, $\text{non}(\bar{w}) \neq \emptyset$. Two words w and w' are *disjoint* if $\text{con}(\bar{w}) \cap \text{con}(\bar{w}') = \emptyset$.

An identity is written as $w \approx w'$, where w and w' are words. An involution semigroup $S = (S, *)$ satisfies an identity $w \approx w'$ if, for any substitution $\varphi : \mathcal{A} \rightarrow S$, the elements $w\varphi$ and $w'\varphi$ of S are equal. An identity $w \approx w'$ is *deducible* from a set Σ of identities, written $\Sigma \vdash w \approx w'$, if every involution semigroup that satisfies all identities in Σ , satisfies $w \approx w'$ as well. Two sets of identities Σ_1 and Σ_2 are *equivalent*, written $\Sigma_1 \sim \Sigma_2$, if the deductions $\Sigma_1 \vdash \Sigma_2$ and $\Sigma_2 \vdash \Sigma_1$ hold. The set of all identities satisfied by an involution semigroup S is denoted by $\text{id}(S)$. A subset Σ of $\text{id}(S)$ is a *basis* for S if $\Sigma \vdash \text{id}(S)$. An involution semigroup is *finitely based* if it possesses a finite basis.

Lemma 2.1 ([10, Lemma 2.1]). *If $w \approx w' \in \text{id}(L)$, then $\text{sim}(\bar{w}) = \text{sim}(\bar{w}')$ and $\text{non}(\bar{w}) = \text{non}(\bar{w}')$.*

Lemma 2.2 ([10, Lemma 2.2]). *For any $w, w' \in \mathcal{A}^+$, the involution semigroup $Z_n^{\mathbb{R}}$ satisfies the identity $w \approx w'$ if and only if $\text{occ}(x, \bar{w}) \equiv \text{occ}(x, \bar{w}') \pmod{n}$ for all $x \in \mathcal{A}$.*

3 An explicit basis for $L \times Z_n^{\mathbb{R}}$

If $n \in \{1, 2\}$, then $\text{sq}(n) \setminus \{1\} = \emptyset$, so that $L \times Z_n^{\mathbb{R}} = L \times Z_n^1$ is finitely based by Theorem 1.2. If $n \geq 3$, then $n - 1 \in \text{sq}(n) \setminus \{1\}$, so that $\text{sq}(n) \setminus \{1\} \neq \emptyset$. Therefore, for the remainder of the article, assume that $n \geq 3$ and $\mathbb{R} \in \text{sq}(n) \setminus \{1\}$ are fixed. Furthermore, the axioms

$$(xy)z \approx x(yz), \quad (x^*)^* \approx x, \quad (xy)^* \approx y^*x^* \tag{inv}$$

of involution semigroups are assumed to hold in all deductions. In other words, for any set Σ of identities and any identity $w \approx w'$, the deduction $\{(inv)\} \cup \Sigma \vdash w \approx w'$ is simply written as $\Sigma \vdash w \approx w'$.

Theorem 3.1. *The identities*

$$x^{n+2} \approx x^2, \quad x^{n+1}yx \approx xyx, \quad xyx^{n+1} \approx xyx, \tag{3.1a}$$

$$x^2yx \approx xyx^2, \tag{3.1b}$$

$$xyxzx \approx xzxyx, \tag{3.1c}$$

$$(x^n)^* \approx x^n, \tag{3.1d}$$

$$x^*h_1x^* \approx x^{\mathbb{R}}h_1x^{\mathbb{R}}, \tag{3.1e}$$

$$x^*h_1xh_2x \approx x^{\mathbb{R}}h_1xh_2x, \tag{3.1f}$$

$$xh_1x^*h_2x \approx xh_1x^{\mathbb{R}}h_2x, \tag{3.1g}$$

$$xh_1xh_2x^* \approx xh_1xh_2x^{\mathbb{R}}, \tag{3.1h}$$

$$x^*h_1y^{\otimes 1}h_2x^{\otimes 2}h_3y^{\otimes 3} \approx x^{\mathbb{R}}h_1y^{\otimes 1}h_2x^{\otimes 2}h_3y^{\otimes 3}, \tag{3.1i}$$

$$x^{\otimes 1}h_1y^*h_2x^{\otimes 2}h_3y^{\otimes 3} \approx x^{\otimes 1}h_1y^{\mathbb{R}}h_2x^{\otimes 2}h_3y^{\otimes 3}, \tag{3.1j}$$

$$x^{\otimes 1}h_1y^{\otimes 2}h_2x^*h_3y^{\otimes 3} \approx x^{\otimes 1}h_1y^{\otimes 2}h_2x^{\mathbb{R}}h_3y^{\otimes 3}, \tag{3.1k}$$

$$x^{\otimes 1}h_1y^{\otimes 2}h_2x^{\otimes 3}h_3y^* \approx x^{\otimes 1}h_1y^{\otimes 2}h_2x^{\otimes 3}h_3y^{\mathbb{R}}, \tag{3.1l}$$

$$(h_1xyh_2)^{n+1} \approx (h_1yxh_2)^{n+1}, \tag{3.1m}$$

$$x^p h_1 y^q h_2 x h_3 y \approx y^q h_1 x^p h_2 y h_3 x, \quad 1 \leq p, q \leq n, \tag{3.1n}$$

$$x h_1 y h_2 x h_3 y \approx x y^n h_1 y h_2 x h_3 y, \tag{3.1o}$$

$$x h_1 y h_2 x h_3 y \approx y^n x h_1 y h_2 x h_3 y, \tag{3.1p}$$

$$xh_1yh_2xh_3y \approx xh_1x^nh_2xh_3y, \tag{3.1q}$$

$$xh_1yh_2xh_3y \approx xh_1yx^nh_2xh_3y, \tag{3.1r}$$

$$xh_1yh_2xh_3y \approx xh_1yh_2xy^nh_3y, \tag{3.1s}$$

$$xh_1yh_2xh_3y \approx xh_1yh_2y^nxh_3y, \tag{3.1t}$$

$$xh_1yh_2xh_3y \approx xh_1yh_2xh_3x^ny, \tag{3.1u}$$

$$xh_1yh_2xh_3y \approx xh_1yh_2xh_3yx^ny, \tag{3.1v}$$

$$x\left(\prod_{i=1}^m(y_ih_iy_i^*)\right)x^* \approx x\left(\prod_{i=m}^1(y_ih_iy_i^*)\right)x^*, \quad m \geq 2, \tag{3.1w}$$

where $h_i \in \{\emptyset, h_i\}$ and $\otimes_i \in \{1, *\}$ constitute a basis for the involution semigroup $L \times Z_n^R$.

Lemma 3.2. *The involution semigroup $L \times Z_n^R$ satisfies identities (3.1).*

Proof. It follows from [10, Lemma 3.2] that the involution semigroup L satisfies identities (3.1). Since the group Z_n^R satisfies the identities $x^n \approx 1$, $x^* \approx x^R$, and $xy \approx yx$, it is routinely shown that it also satisfies identities (3.1). □

Some restrictions on the identities satisfied by the involution semigroup $L \times Z_n^R$ are established in Sections 4 and 5. The proof of Theorem 3.1 is then given in Section 6. The basis (3.1) is shown in Section 7 to contain an infinite irredundant basis for $L \times Z_n^R$, hence completing the proof of Theorem 1.3.

4 Connected identities and sandwich identities

Recall that two words w_1 and w_2 are *disjoint* if $\text{con}(\overline{w_1}) \cap \text{con}(\overline{w_2}) = \emptyset$. A non-simple word is *connected* if it cannot be decomposed into a product of two disjoint nonempty words. In other words, a non-simple word w is connected if whenever $w = w_1w_2$ for some $w_1, w_2 \in (\mathcal{A} \cup \mathcal{A}^*)^+$, then $\text{con}(\overline{w_1}) \cap \text{con}(\overline{w_2}) \neq \emptyset$. An identity $w \approx w'$ is *connected* if the words w and w' are connected. It is shown in Section 4.1 that the involution semigroup $L \times Z_n^R$ possesses a basis that consists entirely of connected identities. In Section 4.2, a special kind of connected identities, called *sandwich identities*, is introduced. It is shown in Lemma 4.4 that the involution semigroup $L \times Z_n^R$ possesses a basis that consists of (3.1) and sandwich identities.

The results in the present section, together with those from Section 5, are required in Section 6 in the proof of Theorem 3.1.

4.1 Connected identities

Lemma 4.1. *Let $w \approx w' \in \text{id}(L \times Z_n^R)$. Suppose that $w = w_1w_2$ for some disjoint words $w_1, w_2 \in (\mathcal{A} \cup \mathcal{A}^*)^+$. Then, $w' = w'_1w'_2$ for some disjoint words $w'_1, w'_2 \in (\mathcal{A} \cup \mathcal{A}^*)^+$ such that $w_1 \approx w'_1, w_2 \approx w'_2 \in \text{id}(L \times Z_n^R)$. Consequently, w is connected if and only if w' is connected.*

Proof. This result has been established for the case $\mathbb{R} = 1$ ([10, Lemma 4.1]), but its proof is independent of the value of \mathbb{R} . Hence, the present lemma holds. □

Any word $u \in (\mathcal{A} \cup \mathcal{A}^*)^+$ can be decomposed as

$$u = \prod_{i=1}^m p_i,$$

where $p_1, p_2, \dots, p_m \in (\mathcal{A} \cup \mathcal{A}^*)^+$ are pairwise disjoint words each of which is either a singleton or a connected word. It is easily seen that this decomposition of u , called the *natural decomposition* of u , is unique.

Lemma 4.2. *The involution semigroup $L \times Z_n^R$ possesses a basis that consists entirely of connected identities.*

Proof. Let $\mathbf{u} \approx \mathbf{u}'$ be any identity from a basis Σ for $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$. Suppose that $\mathbf{u} = \prod_{i=1}^m \mathbf{p}_i$ is the natural decomposition of \mathbf{u} . Then, it follows from Lemma 4.1 that \mathbf{u}' can be decomposed as $\mathbf{u}' = \prod_{i=1}^m \mathbf{p}'_i$ with $\mathbf{p}_i \approx \mathbf{p}'_i \in \text{id}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ for all i .

Case 1: The word \mathbf{p}_i is a singleton. Since $\mathbf{p}_i \approx \mathbf{p}'_i \in \text{id}(\mathbf{L})$, it follows from Lemma 2.1 that $\mathbf{p}_i, \mathbf{p}'_i \in \{x, x^*\}$ for some $x \in \mathcal{A}$. But $\mathbf{p}_i \approx \mathbf{p}'_i \in \text{id}(\mathbf{Z}_n^{\mathbb{R}})$ and $x \approx x^* \notin \text{id}(\mathbf{Z}_n^{\mathbb{R}})$ imply that $\mathbf{p}_i = \mathbf{p}'_i$.

Case 2: The word \mathbf{p}_i is connected. Then, \mathbf{p}'_i is connected by Lemma 4.1.

Therefore, for each i , the identity $\mathbf{p}_i \approx \mathbf{p}'_i$ is either trivial or connected. Consequently, when the identity $\mathbf{u} \approx \mathbf{u}'$ in Σ is replaced by the connected identities from $\{\mathbf{p}_i \approx \mathbf{p}'_i : 1 \leq i \leq m\}$, the resulting set remains a basis for $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$. □

4.2 Sandwich identities

Let $<$ be a total order on the alphabet \mathcal{A} . For any nonempty finite subset \mathcal{X} of \mathcal{A} , write $\mathcal{X} = \{x_1 < x_2 < \dots < x_r\}$ to indicate that x_1, x_2, \dots, x_r are all the letters of \mathcal{X} with $x_1 < x_2 < \dots < x_r$. For such a set \mathcal{X} , define

$$\mathcal{X}^{\boxplus} = \{x_1^{c_1} x_2^{c_2} \dots x_r^{c_r} : c_1, c_2, \dots, c_r \in \{1, 2, \dots, n\}\}.$$

The shortest word in \mathcal{X}^{\boxplus} is

$$\vec{\mathcal{X}} = x_1 x_2 \dots x_r.$$

For any word $\mathbf{w} \in (\mathcal{A} \cup \mathcal{A}^*)^+$ such that $\text{con}(\bar{\mathbf{w}}) = \{x_1 < x_2 < \dots < x_r\}$, define $\min(\mathbf{w}) = x_1$.

A connected word \mathbf{s} is called a *sandwich* if one of the following holds.

- (S.1) $\mathbf{s} = \mathbf{x}\mathbf{u}\mathbf{x}^*$ for some $x \in \mathcal{A}$ and $\mathbf{u} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $x \notin \text{con}(\bar{\mathbf{u}})$,
- (S.2) $\mathbf{s} = \mathbf{x}^*\mathbf{u}\mathbf{x}$ for some $x \in \mathcal{A}$ and $\mathbf{u} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $x \notin \text{con}(\bar{\mathbf{u}})$,
- (S.3) $\mathbf{s} = \mathbf{x} \prod_{i=1}^{\ell} (\mathbf{u}_i \vec{\mathcal{X}})$ for some $\ell \geq 1$, finite nonempty $\mathcal{X} \subseteq \mathcal{A}$, $\mathbf{x} \in \mathcal{X}^{\boxplus}$, and $\mathbf{u}_i \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that
 - (a) $\vec{\mathcal{X}}, \mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{\ell}$ are pairwise disjoint and
 - (b) if $\ell \geq 2$, then $\mathbf{u}_i \neq \emptyset$ for all i and $\min(\mathbf{u}_1) < \min(\mathbf{u}_2) < \dots < \min(\mathbf{u}_{\ell})$.

Specifically, for any $k \in \{1, 2, 3\}$, a sandwich from (S.k) is said to be of *type* (S.k). The *level* of the sandwich in (S.3) is the number ℓ , while the *level* of any sandwich in (S.1) and (S.2) is defined to be one.

Remark 4.3. In (S.3), due to (b), the only case in which any of $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_{\ell}$ can be empty is when $\ell = 1$ and $\mathbf{u}_1 = \emptyset$, that is, $\mathbf{s} = \mathbf{x}\vec{\mathcal{X}}$.

An identity $\mathbf{s} \approx \mathbf{s}'$ is a *sandwich identity* if the words \mathbf{s} and \mathbf{s}' are sandwiches. Denote by

$$\text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$$

the set of all sandwich identities satisfied by the involution semigroup $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$.

Lemma 4.4. *The set $\{(3.1)\} \cup \text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ constitutes a basis for the involution semigroup $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$.*

Recall from Lemma 4.2 that the set $\text{id}_{\mathbf{C}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ of all connected identities from $\text{id}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ constitutes a basis for $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$. As shown in Lemma 3.2, the involution semigroup $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$ satisfies identities (3.1), so that the set $\{(3.1)\} \cup \text{id}_{\mathbf{C}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ is also a basis for $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$. It is shown in Lemma 4.7 that identities (3.1) can be used to convert any connected word into a sandwich. Lemma 4.4 thus follows.

Lemma 4.5 ([10, Lemma 4.5]). *The following identities are deducible from (3.1):*

$$\left(\prod_{i=1}^r x_i^{c_i}\right) \mathbf{h} \left(\prod_{i=1}^r x_i\right)^2 \approx \left(\prod_{i=1}^r x_i^{c_i+1}\right) \mathbf{h} \left(\prod_{i=1}^r x_i\right), \quad c_i \geq 1, r \geq 1, \tag{4.1a}$$

$$\left(\prod_{i=1}^r x_i^{c_i}\right) \left(\prod_{i=1}^r x_i\right) \mathbf{h} \left(\prod_{i=1}^r x_i\right) \approx \left(\prod_{i=1}^r x_i^{c_i+1}\right) \mathbf{h} \left(\prod_{i=1}^r x_i\right), \quad c_i \geq 1, r \geq 1, \tag{4.1b}$$

where $\mathbf{h} \in \{\emptyset, h\}$.

Lemma 4.6. *Suppose that w is any connected word. Then, identities (3.1) can be used to convert w into some word w' such that $\triangleright \bar{w}' = \bar{w}' \triangleleft$.*

Proof. Suppose that $\triangleright \bar{w} \neq \bar{w} \triangleleft$. Then, since the word w is connected, there exist distinct letters $x_1, \dots, x_r \in \mathcal{A}$ occurring in w in the overlapping manner

$$w = x_1^{\otimes_1} a_1 x_2^{\otimes_2} a_2 x_1^{\otimes'_1} b_1 x_3^{\otimes_3} a_3 x_2^{\otimes'_2} b_2 x_4^{\otimes_4} a_4 x_3^{\otimes'_3} b_3 \cdots x_r^{\otimes_r} a_r x_{r-1}^{\otimes'_{r-1}} b_{r-1} x_r^{\otimes'_r},$$

where $a_i, b_i \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ and $\otimes_i, \otimes'_i \in \{1, *\}$. (Note that a_i follows $x_i^{\otimes_i}$ while b_i follows $x_i^{\otimes'_i}$.) Then, identities {(3.1b), (3.1i), (3.1j), (3.1k), (3.1l)} can be used to convert w into

$$x_1^{p_1} a_1 x_2^{p_2} a_2 x_1 b_1 x_3^{p_3} a_3 x_2 b_2 x_4^{p_4} a_4 x_3 b_3 \cdots x_r^{p_r} a_r x_{r-1} b_{r-1} x_r,$$

where $p_i, q_i \geq 1$. Hence,

$$\begin{aligned} w &\stackrel{(3.1)}{\approx} x_1^{p_1} a_1 x_2^{p_2} a_2 x_1 b_1 x_3^{p_3} a_3 x_2 b_2 x_4^{p_4} a_4 x_3 b_3 \cdots x_r^{p_r} a_r x_{r-1} b_{r-1} x_r \\ &\stackrel{(3.1v)}{\approx} x_1^{p_1} a_1 x_2^{p_2} a_2 x_1 b_1 x_3^{p_3} a_3 x_2 x_1^n b_2 x_4^{p_4} a_4 x_3 b_3 \cdots x_r^{p_r} a_r x_{r-1} b_{r-1} x_r \\ &\stackrel{(3.1v)}{\approx} x_1^{p_1} a_1 x_2^{p_2} a_2 x_1 b_1 x_3^{p_3} a_3 x_2 x_1^n b_2 x_4^{p_4} a_4 x_3 x_1^n b_3 \cdots x_r^{p_r} a_r x_{r-1} b_{r-1} x_r \\ &\vdots \\ &\stackrel{(3.1v)}{\approx} x_1^{p_1} a_1 x_2^{p_2} a_2 x_1 b_1 x_3^{p_3} a_3 x_2 x_1^n b_2 x_4^{p_4} a_4 x_3 x_1^n b_3 \cdots x_r^{p_r} a_r x_{r-1} x_1^n b_{r-1} x_r \\ &\stackrel{(3.1v)}{\approx} x_1^{p_1} a_1 x_2^{p_2} a_2 x_1 b_1 x_3^{p_3} a_3 x_2 x_1^n b_2 x_4^{p_4} a_4 x_3 x_1^n b_3 \cdots x_r^{p_r} a_r x_{r-1} x_1^n b_{r-1} x_r x_1^n \\ &= w', \end{aligned}$$

where $\triangleright \bar{w}' = x_1 = \bar{w}' \triangleleft$. □

Lemma 4.7. *Suppose that w is any connected word. Then, identities (3.1) can be used to convert w into some sandwich s with $\text{sim}(\bar{w}) = \text{sim}(\bar{s})$ and $\text{non}(\bar{w}) = \text{non}(\bar{s})$.*

Proof. By Lemma 4.5, it suffices to convert the word w , using identities (3.1) and (4.1), into some sandwich s with $\text{sim}(\bar{w}) = \text{sim}(\bar{s})$ and $\text{non}(\bar{w}) = \text{non}(\bar{s})$. By Lemma 4.6, it can be assumed that $x_1 = \triangleright \bar{w} = \bar{w} \triangleleft$. Then, w can be written as

$$w = x_1^{\otimes_0} \prod_{i=1}^{m_1} (w_{1,i} x_1^{\otimes_i}),$$

where $m_1 \geq 1$, $\otimes_i \in \{1, *\}$, and $w_{1,i} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ with $x_1 \notin \text{con}(\bar{w}_{1,i})$ for all i . If $m_1 = 1$, then w is either $x_1 w_{1,1} x_1$, $x_1 w_{1,1} x_1^*$, $x_1^* w_{1,1} x_1$, or $x_1^* w_{1,1} x_1^*$; the first three words are sandwiches while identities (3.1) can be used to convert the fourth word into a sandwich of type (S.3), that is,

$$x_1^* w_{1,1} x_1^* \stackrel{(3.1e)}{\approx} x_1^R w_{1,1} x_1^R \stackrel{(3.1b)}{\approx} x_1^{2R-1} w_{1,1} x_1 \stackrel{(3.1a)}{\approx} x_1^c w_{1,1} x_1,$$

where $c \in \{1, 2, \dots, n\}$ is such that $c \equiv 2R - 1 \pmod{n}$. Therefore, assume that $m_1 \geq 2$, so that identities (3.1e)–(3.1h) can be used to replace any x_1^* by x_1^R .

Suppose that $w_{1,i}$ and $w_{1,j}$ are not disjoint with $i \neq j$, say $x_2 \in \text{con}(\bar{w}_{1,i}) \cap \text{con}(\bar{w}_{1,j})$. Then,

$$w = x_1 h_1 x_2^{\otimes'_1} h_2 x_1 h_3 x_2^{\otimes'_2} h_4 x_1$$

for some $h_i \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ and $\otimes'_i \in \{1, *\}$. Identities (3.1i)–(3.1l) can first be used to replace any x_2^* by x_2^R and identities (3.1o)–(3.1v) can then be used to perform the replacement $(x_1, x_2) \mapsto (x_1 x_2^n, x_1^n x_2)$. The resulting word is of the form

$$x_1^{c_{0,1}} x_2^{c_{0,2}} \prod_{i=1}^{m_2} (w_{2,i} x_1^{c_{i,1}} x_2^{c_{i,2}}),$$

where $m_2 \geq 2$, $c_{i,j} \geq 1$, and $w_{2,i} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ with $x_1, x_2 \notin \text{con}(\overline{w_{2,i}})$ for all i . Similarly, if $w_{2,i}$ and $w_{2,j}$ are not disjoint with $i \neq j$, say $x_3 \in \text{con}(\overline{w_{2,i}}) \cap \text{con}(\overline{w_{2,j}})$, then identities (3.1i)–(3.1l) can first be used to replace any x_3^* by x_3^R and identities (3.1o)–(3.1v) can then be used to perform the replacement $(x_2^{c_{i,2}}, x_3) \mapsto (x_2^{c_{i,2}} x_3^n, x_1^n x_2^n x_3)$. The resulting word is of the form

$$x_1^{c_{0,1}} x_2^{c_{0,2}} x_3^{c_{0,3}} \prod_{i=1}^{m_3} (w_{3,i} x_1^{c_{i,1}} x_2^{c_{i,2}} x_3^{c_{i,3}}),$$

where $m_3 \geq 2$, $c_{i,j} \geq 1$, and $w_{3,i} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ with $x_1, x_2, x_3 \notin \text{con}(\overline{w_{3,i}})$ for all i . This can be repeated until a word of the form

$$x_1^{c_{0,1}} x_2^{c_{0,2}} \cdots x_r^{c_{0,r}} \prod_{i=1}^{m_r} (w_{r,i} x_1^{c_{i,1}} x_2^{c_{i,2}} \cdots x_r^{c_{i,r}})$$

is obtained, where $m_r \geq 2$, $c_{i,j} \geq 1$, and $w_{r,1}, w_{r,2}, \dots, w_{r,m_r} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ are pairwise disjoint words with $x_1, x_2, \dots, x_r \notin \text{con}(\overline{w_{r,i}})$ for all i . It is easily seen that identities (3.1a) and (3.1b) can be used to convert this word into

$$w' = x_1^{c_1} x_2^{c_2} \cdots x_r^{c_r} \prod_{i=1}^{m_r} (w_{r,i} x_1 x_2 \cdots x_r),$$

where $1 \leq c_i \leq n$. Let π denote the permutation on $\{1, 2, \dots, r\}$ such that

$$x_{1\pi} < x_{2\pi} < \cdots < x_{r\pi}.$$

Then,

$$\begin{aligned} w' &\stackrel{(3.1a)}{\approx} x_1^{c_1} x_2^{c_2} \cdots x_r^{c_r} w_{r,1} x_1 x_2 \cdots x_r \prod_{i=2}^{m_r} (w_{r,i} (x_1 x_2 \cdots x_r)^{n+1}) \\ &\stackrel{(3.1n)}{\approx} x_{1\pi}^{c_{1\pi}} x_{2\pi}^{c_{2\pi}} \cdots x_{r\pi}^{c_{r\pi}} w_{r,1} x_{1\pi} x_{2\pi} \cdots x_{r\pi} \prod_{i=2}^{m_r} (w_{r,i} (x_{1\pi} x_{2\pi} \cdots x_{r\pi})^{n+1}) \\ &\stackrel{(3.1m)}{\approx} x_{1\pi}^{c_{1\pi}} x_{2\pi}^{c_{2\pi}} \cdots x_{r\pi}^{c_{r\pi}} w_{r,1} x_{1\pi} x_{2\pi} \cdots x_{r\pi} \prod_{i=2}^{m_r} (w_{r,i} (x_{1\pi} x_{2\pi} \cdots x_{r\pi})^{n+1}) \\ &\stackrel{(3.1a)}{\approx} x_{1\pi}^{c_{1\pi}} x_{2\pi}^{c_{2\pi}} \cdots x_{r\pi}^{c_{r\pi}} \prod_{i=1}^{m_r} (w_{r,i} x_{1\pi} x_{2\pi} \cdots x_{r\pi}). \end{aligned}$$

In summary, identities (3.1) can be used to convert w into a word of the form

$$s = x \prod_{i=1}^{\ell} (u_i \vec{\mathcal{X}}),$$

where $\ell \geq 1$, $\mathcal{X} = \{x_1 < x_2 < \cdots < x_r\} \subseteq \mathcal{A}$, $x = x_1^{c_1} x_2^{c_2} \cdots x_r^{c_r} \in \mathcal{X}^{\boxplus}$, and $u_i \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ are such that $\vec{\mathcal{X}}, u_1, u_2, \dots, u_\ell$ are pairwise disjoint. Let c'_i be the number in $\{1, 2, \dots, n\}$ such that $c_i + 1 \equiv c'_i \pmod{n}$. If $u_1 = \emptyset$, then

$$s = x_1^{c_1} x_2^{c_2} \cdots x_r^{c_r} \vec{\mathcal{X}} u_2 \vec{\mathcal{X}} \prod_{i=3}^{\ell} (u_i \vec{\mathcal{X}}) \stackrel{(4.1b)}{\approx} x_1^{c_1+1} x_2^{c_2+1} \cdots x_r^{c_r+1} u_2 \vec{\mathcal{X}} \prod_{i=3}^{\ell} (u_i \vec{\mathcal{X}}) \stackrel{(3.1a)}{\approx} x_1^{c'_1} x_2^{c'_2} \cdots x_r^{c'_r} u_2 \vec{\mathcal{X}} \prod_{i=3}^{\ell} (u_i \vec{\mathcal{X}})$$

and if $u_j = \emptyset$ for some $j \in \{2, 3, \dots, \ell\}$, then

$$\begin{aligned} s &= x_1^{c_1} x_2^{c_2} \cdots x_r^{c_r} \left(\prod_{i=1}^{j-1} (u_i \vec{\mathcal{X}}) \right) \vec{\mathcal{X}} \left(\prod_{i=j+1}^{\ell} (u_i \vec{\mathcal{X}}) \right) \\ &\stackrel{(4.1a)}{\approx} x_1^{c_1+1} x_2^{c_2+1} \cdots x_r^{c_r+1} \left(\prod_{i=1}^{j-1} (u_i \vec{\mathcal{X}}) \right) \left(\prod_{i=j+1}^{\ell} (u_i \vec{\mathcal{X}}) \right) \\ &\stackrel{(3.1a)}{\approx} x_1^{c'_1} x_2^{c'_2} \cdots x_r^{c'_r} \left(\prod_{i=1}^{j-1} (u_i \vec{\mathcal{X}}) \right) \left(\prod_{i=j+1}^{\ell} (u_i \vec{\mathcal{X}}) \right). \end{aligned}$$

Hence, for any i , if the factor u_i is empty, then the $\vec{\mathcal{X}}$ that follows it can be “combined” with the prefix x . Therefore, it can further be assumed that either

- (A) $s = x \vec{\mathcal{X}}$ or
- (B) $u_i \neq \emptyset$ for all i .

If (A) holds, then the word s is a sandwich of type (S.3). Hence, assume that (B) holds. It remains to show that if $\ell \geq 2$, then identities (3.1) can be used to rearrange the factors $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_\ell$, so that s satisfies (S.3)(b) and is a sandwich of type (S.3). To interchange \mathbf{u}_j and \mathbf{u}_{j+1} for any $j \geq 2$, identity (3.1c) can clearly be used. To interchange \mathbf{u}_1 and \mathbf{u}_2 ,

$$\begin{aligned} s &\stackrel{(3.1a)}{\approx} \mathbf{x} \mathbf{u}_1 \vec{\mathcal{X}} \mathbf{u}_2 \vec{\mathcal{X}} \vec{\mathcal{X}} \vec{\mathcal{X}}^{n-1} \prod_{i=3}^{\ell} (\mathbf{u}_i \vec{\mathcal{X}}) \\ &\stackrel{(3.1b)}{\approx} \vec{\mathcal{X}} \mathbf{u}_1 \vec{\mathcal{X}} \mathbf{u}_2 \vec{\mathcal{X}} \mathbf{x} \vec{\mathcal{X}}^{n-1} \prod_{i=3}^{\ell} (\mathbf{u}_i \vec{\mathcal{X}}) \\ &\stackrel{(3.1c)}{\approx} \vec{\mathcal{X}} \mathbf{u}_2 \vec{\mathcal{X}} \mathbf{u}_1 \vec{\mathcal{X}} \mathbf{x} \vec{\mathcal{X}}^{n-1} \prod_{i=3}^{\ell} (\mathbf{u}_i \vec{\mathcal{X}}) \\ &\stackrel{(3.1b)}{\approx} \mathbf{x} \mathbf{u}_2 \vec{\mathcal{X}} \mathbf{u}_1 \vec{\mathcal{X}} \vec{\mathcal{X}} \vec{\mathcal{X}}^{n-1} \prod_{i=3}^{\ell} (\mathbf{u}_i \vec{\mathcal{X}}) \\ &\stackrel{(3.1a)}{\approx} \mathbf{x} \mathbf{u}_2 \vec{\mathcal{X}} \mathbf{u}_1 \vec{\mathcal{X}} \prod_{i=3}^{\ell} (\mathbf{u}_i \vec{\mathcal{X}}). \end{aligned}$$

Throughout this proof, identities (3.1) have been used to convert the word w into a sandwich s . Therefore, $w \approx s \in \text{id}(\mathbf{L})$ by Lemma 3.2, so that $\text{sim}(\overline{w}) = \text{sim}(\overline{s})$ and $\text{non}(\overline{w}) = \text{non}(\overline{s})$ by Lemma 2.1. □

5 Restrictions on sandwich identities

The present section establishes some properties of sandwich identities satisfied by the involution semigroup $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$. In Section 5.1, it is shown that any two sandwiches that form such an identity must share the same type and level. In Section 5.2, *refined identities* are introduced; these are identities formed by certain sandwiches of level one. It is shown in Lemma 5.8 that refined identities satisfied by the involution semigroup $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$ are of very specific form.

The results established in the present section are required in Section 6 in the proof of Theorem 3.1.

5.1 Type of sandwiches forming sandwich identities

Lemma 5.1. *Suppose that $s \approx s' \in \text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$, where $s = \mathbf{x} \mathbf{u} \mathbf{x}^*$ is the sandwich in (S.1). Then, $s' = \mathbf{x} \mathbf{u}' \mathbf{x}^*$ for some $\mathbf{u}' \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ with $\mathbf{x} \notin \text{con}(\overline{\mathbf{u}}) = \text{con}(\overline{\mathbf{u}'})$. Consequently, s is of type (S.1) if and only if s' is of type (S.1).*

Proof. Let $\varphi : \mathcal{A} \rightarrow L$ denote the substitution given by

$$z\varphi = \begin{cases} fe & \text{if } z = x, \\ e & \text{otherwise.} \end{cases}$$

Then, $s\varphi = fef = s'\varphi$. It follows that

$$s'\varphi = fe \cdot e \cdot e \cdots e \cdot ef,$$

whence $s' = \mathbf{x} \mathbf{u}' \mathbf{x}^*$ for some $\mathbf{u}' \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ with $\mathbf{x} \notin \text{con}(\overline{\mathbf{u}'})$. Furthermore, $\text{con}(\overline{s}) = \text{con}(\overline{s'})$ by Lemma 2.1, so that $\mathbf{x} \notin \text{con}(\overline{\mathbf{u}}) = \text{con}(\overline{\mathbf{u}'})$. □

Lemma 5.2. *Suppose that $s \approx s' \in \text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$, where $s = \mathbf{x}^* \mathbf{u} \mathbf{x}$ is the sandwich in (S.2). Then, $s' = \mathbf{x}^* \mathbf{u}' \mathbf{x}$ for some $\mathbf{u}' \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ with $\mathbf{x} \notin \text{con}(\overline{\mathbf{u}}) = \text{con}(\overline{\mathbf{u}'})$. Consequently, s is of type (S.2) if and only if s' is of type (S.2).*

Proof. This is symmetric to Lemma 5.1. □

Lemma 5.3. Suppose that $s \approx s' \in \text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$, where

$$\mathbf{s} = \mathbf{x} \prod_{i=1}^{\ell} (\mathbf{u}_i \vec{\mathcal{X}})$$

is the sandwich in (S.3). Then,

$$\mathbf{s}' = \mathbf{x} \prod_{i=1}^{\ell} (\mathbf{u}'_i \vec{\mathcal{X}})$$

for some $\mathbf{u}'_1, \mathbf{u}'_2, \dots, \mathbf{u}'_{\ell} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $\text{con}(\overline{\mathbf{u}}_i) = \text{con}(\overline{\mathbf{u}'_i})$ for all i . Consequently, \mathbf{s} is of type (S.3) if and only if \mathbf{s}' is of type (S.3).

Proof. Following the proof of [10, Lemma 5.3], there holds

$$\mathbf{s}' = \mathbf{y} \prod_{i=1}^{\ell} (\mathbf{u}'_i \vec{\mathcal{X}})$$

for some $\mathbf{y} \in \mathcal{X}^{\text{th}}$ and $\mathbf{u}'_1, \mathbf{u}'_2, \dots, \mathbf{u}'_{\ell} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $\text{con}(\overline{\mathbf{u}}_i) = \text{con}(\overline{\mathbf{u}'_i})$ for all i . Since

$$\mathcal{X} \cap \text{con}(\overline{\mathbf{u}}_1 \cdot \overline{\mathbf{u}}_2 \cdots \overline{\mathbf{u}}_{\ell}) = \emptyset$$

by (S.3)(a) and $s \approx s' \in \text{id}(\mathbf{Z}_n^{\mathbb{R}})$, where \mathbf{Z}_n has a unit element, $\mathbf{Z}_n^{\mathbb{R}}$ satisfies the identity $\mathbf{x} \vec{\mathcal{X}}^{\ell} \approx \mathbf{y} \vec{\mathcal{X}}^{\ell}$ obtained from $s \approx s'$ by retaining the letters in \mathcal{X} . Since $\mathbf{x} \vec{\mathcal{X}}^{\ell}, \mathbf{y} \vec{\mathcal{X}}^{\ell} \in \mathcal{A}^+$, it follows from Lemma 2.2 that $\mathbf{x} = \mathbf{y}$. \square

Lemma 5.4. Suppose that $s \approx s' \in \text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$, where

$$\mathbf{s} = \mathbf{x} \prod_{i=1}^{\ell} (\mathbf{u}_i \vec{\mathcal{X}}) \quad \text{and} \quad \mathbf{s}' = \mathbf{x} \prod_{i=1}^{\ell} (\mathbf{u}'_i \vec{\mathcal{X}})$$

are the sandwiches of type (S.3) in Lemma 5.3. Then,

$$\{(3.1), \mathbf{s} \approx \mathbf{s}'\} \sim \{(3.1)\} \cup \{\mathbf{x} \mathbf{u}_i \mathbf{x} \approx \mathbf{x} \mathbf{u}'_i \mathbf{x} : 1 \leq i \leq \ell\}.$$

Proof. For each $i \in \{1, 2, \dots, \ell\}$, let $\varphi_i : \mathcal{A} \rightarrow \mathcal{A}^+$ denote the substitution given by

$$z\varphi_i = \begin{cases} z & \text{if } z \in \text{con}(\overline{\mathbf{u}}_i) = \text{con}(\overline{\mathbf{u}'_i}), \\ \mathbf{x}^n & \text{otherwise.} \end{cases}$$

Note that for any $\mathbf{w} \in (\mathcal{A} \cup \mathcal{A}^*)^+$ such that \mathbf{w} and \mathbf{u}_i are disjoint, the image $\mathbf{w}\varphi_i$ belongs to $\{\mathbf{x}^n, (\mathbf{x}^n)^*\}^+$. Therefore, identity (3.1d) can be used to convert $\mathbf{w}\varphi_i$ into the plain word $\overline{\mathbf{w}\varphi_i}$ in $\{\mathbf{x}^n\}^+$. Hence, by (S.3)(a),

$$\begin{aligned} \mathbf{x}(\mathbf{s}\varphi_i)\mathbf{x} &= \mathbf{x}((\mathbf{x}\mathbf{u}_1 \vec{\mathcal{X}} \cdots \mathbf{u}_{i-1} \vec{\mathcal{X}})\varphi_i)\mathbf{u}_i((\vec{\mathcal{X}}\mathbf{u}_{i+1} \vec{\mathcal{X}} \cdots \mathbf{u}_{\ell} \vec{\mathcal{X}})\varphi_i)\mathbf{x} \\ &\stackrel{(3.1d)}{\approx} \mathbf{x}(\overline{(\mathbf{x}\mathbf{u}_1 \vec{\mathcal{X}} \cdots \mathbf{u}_{i-1} \vec{\mathcal{X}})\varphi_i})\mathbf{u}_i(\overline{(\vec{\mathcal{X}}\mathbf{u}_{i+1} \vec{\mathcal{X}} \cdots \mathbf{u}_{\ell} \vec{\mathcal{X}})\varphi_i})\mathbf{x} \\ &\stackrel{(3.1a)}{\approx} \mathbf{x}\mathbf{u}_i\mathbf{x}, \end{aligned}$$

so that (3.1) $\vdash \mathbf{x}(\mathbf{s}\varphi_i)\mathbf{x} \approx \mathbf{x}\mathbf{u}_i\mathbf{x}$. Similarly, the deduction (3.1) $\vdash \mathbf{x}(\mathbf{s}'\varphi_i)\mathbf{x} \approx \mathbf{x}\mathbf{u}'_i\mathbf{x}$ holds. Thus, the deduction $\{(3.1), \mathbf{s} \approx \mathbf{s}'\} \vdash \mathbf{x}\mathbf{u}_i\mathbf{x} \approx \mathbf{x}\mathbf{u}'_i\mathbf{x}$ holds. Conversely,

$$\begin{aligned} \mathbf{s} &= \mathbf{x}_1^{c_1} \mathbf{x}_2^{c_2} \cdots \mathbf{x}_r^{c_r} \mathbf{u}_1 \vec{\mathcal{X}} \prod_{i=2}^{\ell} (\mathbf{u}_i \vec{\mathcal{X}}) \\ &\stackrel{(3.1b)}{\approx} \mathbf{x}_1^{c_1} \mathbf{x}_2^{c_2} \cdots \mathbf{x}_r^{c_r} \mathbf{u}_1 \mathbf{x}_r^n \vec{\mathcal{X}} \prod_{i=2}^{\ell} (\mathbf{u}_i \vec{\mathcal{X}}) \\ &\approx \mathbf{x}_1^{c_1} \mathbf{x}_2^{c_2} \cdots \mathbf{x}_r^{c_r} \mathbf{u}'_1 \mathbf{x}_r^n \vec{\mathcal{X}} \prod_{i=2}^{\ell} (\mathbf{u}'_i \vec{\mathcal{X}}) \quad \text{by } \mathbf{x}\mathbf{u}_i\mathbf{x} \approx \mathbf{x}\mathbf{u}'_i\mathbf{x} \\ &\stackrel{(3.1b)}{\approx} \mathbf{s}'. \end{aligned}$$

Therefore, the deduction $\{(3.1)\} \cup \{\mathbf{x}\mathbf{u}_i\mathbf{x} \approx \mathbf{x}\mathbf{u}'_i\mathbf{x} : 1 \leq i \leq \ell\} \vdash \mathbf{s} \approx \mathbf{s}'$ holds. \square

5.2 Refined identities

By Lemmas 5.1–5.3, any identity in $\text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ is formed by a pair of sandwiches that share the same type and level. Therefore, it is unambiguous to define the *type* and *level* of a sandwich identity $s \approx s'$ in $\text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ to be, respectively, the type and level shared by the sandwiches s and s' . The present subsection investigates identities in $\text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ of level one.

Consider a word \mathbf{r} of the form

$$\mathbf{r} = x \left(\prod_{i=1}^k \mathbf{p}_i \right) x^{\otimes}, \tag{5.1}$$

where $k \geq 1$, $x \in \mathcal{A}$, $\otimes \in \{1, *\}$, and $\mathbf{p}_i \in (\mathcal{A} \cup \mathcal{A}^*)^+$ are such that $x, \mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k$ are pairwise disjoint. Note that depending on \otimes , the word \mathbf{r} is a level one sandwich of type (S.1) or (S.3). This sandwich is said to be *refined* if it satisfies both that

- (R.1) each \mathbf{p}_i is either a singleton or a sandwich and
- (R.2) if $\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k$ are all sandwiches with $k \geq 2$, then $\min(\mathbf{p}_1) < \min(\mathbf{p}_k)$.

An identity $\mathbf{r} \approx \mathbf{r}'$ is a *refined identity* if the words \mathbf{r} and \mathbf{r}' are refined sandwiches. Denote by

$$\text{id}_{\text{REF}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$$

the set of all refined identities satisfied by the involution semigroup $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$.

Lemma 5.5. *Suppose that $s \approx s' \in \text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$. Then, there exists some finite subset Σ of $\text{id}_{\text{REF}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ such that the equivalence $\{(3.1), s \approx s'\} \sim \{(3.1)\} \cup \Sigma$ holds. Furthermore, each identity $\mathbf{r} \approx \mathbf{r}'$ in Σ can be chosen so that $|\text{sim}(\bar{\mathbf{r}})| \leq |\text{sim}(\bar{\mathbf{s}})|$ and $|\text{non}(\bar{\mathbf{r}})| \leq |\text{non}(\bar{\mathbf{s}})|$.*

Proof. There are three cases depending on the type of the sandwich identity $s \approx s'$.

Case 1: The word $s \approx s'$ is of type (S.1). Then, by Lemma 5.1,

$$s = x u x^* \quad \text{and} \quad s' = x u' x^*$$

for some $x \in \mathcal{A}$ and $u, u' \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $x \notin \text{con}(\bar{u}) = \text{con}(\bar{u}')$. If $\text{con}(\bar{u}) = \text{con}(\bar{u}') = \emptyset$, then the identity $s \approx s'$ is trivial, so that the result holds with $\Sigma = \emptyset$. Hence, assume that $\text{con}(\bar{u}) = \text{con}(\bar{u}') \neq \emptyset$. In what follows, it is shown that identities (3.1) can be used to convert s into some refined sandwich \mathbf{r} . Similarly, identities (3.1) can be used to convert s' into some refined sandwich \mathbf{r}' . Hence, the equivalence $\{(3.1), s \approx s'\} \sim \{(3.1), \mathbf{r} \approx \mathbf{r}'\}$ holds. Since the deduction $(3.1) \vdash s \approx \mathbf{r}$ holds, it follows from Lemma 3.2 that $s \approx \mathbf{r} \in \text{id}(\mathbf{L})$, whence $|\text{sim}(\bar{\mathbf{r}})| = |\text{sim}(\bar{\mathbf{s}})|$ and $|\text{non}(\bar{\mathbf{r}})| = |\text{non}(\bar{\mathbf{s}})|$ by Lemma 2.1.

Let $u = \prod_{i=1}^k \mathbf{p}_i$ be the natural decomposition of u , so that $x, \mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k$ are pairwise disjoint and each \mathbf{p}_i is either a singleton or a connected word. By Lemma 4.7, identities (3.1) can be used to convert any connected \mathbf{p}_i into some sandwich \mathbf{s}_i with $\text{sim}(\bar{\mathbf{p}}_i) = \text{sim}(\bar{\mathbf{s}}_i)$ and $\text{non}(\bar{\mathbf{p}}_i) = \text{non}(\bar{\mathbf{s}}_i)$. Therefore, it can be assumed that s satisfies (R.1). If s also satisfies (R.2), then s is already refined. Hence, suppose that s does not satisfy (R.2), that is, $\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_k$ are all sandwiches with $k \geq 2$, but $\min(\mathbf{p}_1) \not< \min(\mathbf{p}_k)$. Then, $\min(\mathbf{p}_k) < \min(\mathbf{p}_1)$ because \mathbf{p}_1 and \mathbf{p}_k are disjoint. Let x_1 be the first letter of $\bar{\mathbf{p}}_1$. If \mathbf{p}_i is of type (S.1), then $\mathbf{p}_i = x_1 \mathbf{w} x_1^*$ for some $\mathbf{w} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$. If \mathbf{p}_i is of type (S.2), then $\mathbf{p}_i = x_1^* \mathbf{w} x_1$ for some $\mathbf{w} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$, so that

$$\mathbf{p}_i \stackrel{(\text{inv})}{\approx} x_1^* \mathbf{w} (x_1^*)^*.$$

If \mathbf{p}_i is of type (S.3), then

$$\mathbf{p}_i = x_1^{c_1} x_2^{c_2} \cdots x_r^{c_r} \mathbf{w} x_1 x_2 \cdots x_r$$

for some $\mathbf{w} \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ and $\mathcal{X} = \{x_1 < x_2 < \cdots < x_r\} \subseteq \mathcal{A}$, so that

$$\mathbf{p}_i \stackrel{(3.1a)}{\approx} x_1^n \mathbf{p}_i \stackrel{(3.1v)}{\approx} x_1^n \mathbf{p}_i x_1^n \stackrel{(3.1h)}{\approx} x_1^n \mathbf{p}_i x_1^{n-R} x_1^*.$$

Therefore, regardless of type, there exist $h_i \in \mathcal{A} \cup \mathcal{A}^*$ and $\mathbf{w}_i \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $\mathbf{p}_i \stackrel{(3.1)}{\approx} h_i \mathbf{w}_i h_i^*$. Hence,

$$s = x \left(\prod_{i=1}^k \mathbf{p}_i \right) x^* \stackrel{(3.1)}{\approx} x \left(\prod_{i=1}^k (h_i \mathbf{w}_i h_i^*) \right) x^* \stackrel{(3.1w)}{\approx} x \left(\prod_{i=k}^1 (h_i \mathbf{w}_i h_i^*) \right) x^* \stackrel{(3.1)}{\approx} x \left(\prod_{i=k}^1 \mathbf{p}_i \right) x^*.$$

Since $\min(\mathbf{p}_k) < \min(\mathbf{p}_1)$, the word $\mathbf{r} = x \left(\prod_{i=k}^1 \mathbf{p}_i \right) x^*$ is the required refined sandwich.

Case 2: The word $s \approx s'$ is of type (S.2). Then, by Lemma 5.2,

$$s = x^* u x \quad \text{and} \quad s' = x^* u' x$$

for some $x \in \mathcal{A}$ and $u, u' \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $x \notin \text{con}(\bar{u}) = \text{con}(\bar{u}')$. It is clear that the equivalence $\{(3.1), s \approx s'\} \sim \{(3.1), x u x^* \approx x u' x^*\}$ holds. Since $x u x^* \approx x u' x^*$ is a sandwich identity of type (S.1), the result follows from Case 1.

Case 3: The word $s \approx s'$ is of type (S.3). Then, by Lemma 5.3,

$$s = x \prod_{i=1}^{\ell} (u_i \bar{\mathcal{X}}) \quad \text{and} \quad s' = x \prod_{i=1}^{\ell} (u'_i \bar{\mathcal{X}})$$

for some $\ell \geq 1$, finite nonempty $\bar{\mathcal{X}} \subseteq \mathcal{A}$, $x \in \bar{\mathcal{X}}^{\#}$, and $u_i, u'_i \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $\text{con}(\bar{u}_i) = \text{con}(\bar{u}'_i)$ for each i and $\bar{\mathcal{X}}, u_1, u_2, \dots, u_{\ell}$ are pairwise disjoint. By Lemma 5.4, the equivalence

(A) $\{(3.1), s \approx s'\} \sim \{(3.1)\} \cup \{x u_i x \approx x u'_i x : 1 \leq i \leq \ell\}$

holds. It is easily seen that

(B) $|\text{sim}(\overline{x u_i x})| \leq |\text{sim}(\bar{s})|$ and $|\text{non}(\overline{x u_i x})| \leq |\text{non}(\bar{s})|$ for all i .

For each i , the arguments in Case 1 can be repeated to show that the equivalence

(C) $\{(3.1), x u_i x \approx x u'_i x\} \sim \{(3.1), r_i \approx r'_i\}$

holds for some $r_i \approx r'_i \in \text{id}_{\text{REF}}(\mathbb{L} \times \mathbb{Z}_n^{\mathbb{R}})$ such that

(D) $|\text{sim}(\bar{r}_i)| = |\text{sim}(\overline{x u_i x})|$ and $|\text{non}(\bar{r}_i)| = |\text{non}(\overline{x u_i x})|$.

Hence, the equivalence $\{(3.1), s \approx s'\} \sim \{(3.1)\} \cup \{r_i \approx r'_i : 1 \leq i \leq \ell\}$ holds by (A) and (C), where $|\text{sim}(\bar{r}_i)| \leq |\text{sim}(\bar{s})|$ and $|\text{non}(\bar{r}_i)| \leq |\text{non}(\bar{s})|$ by (B) and (D). □

Lemma 5.6 ([8, Lemma 13]). *Let $w \approx w' \in \text{id}(\mathbb{L})$, where $w, w' \in \mathcal{A}^+$. Suppose that w belongs to*

$$\mathfrak{P}_k^{\uparrow} = \{x^{q_1} y_1^{t_1} y_2^{t_2} \cdots y_k^{t_k} x^{q_2} : q_i \geq 1, t_j \geq 2\}$$

for some $k \geq 2$. Then, w' belongs to either $\mathfrak{P}_k^{\uparrow}$ or

$$\mathfrak{P}_k^{\downarrow} = \{x^{q_1} y_k^{t_k} y_{k-1}^{t_{k-1}} \cdots y_1^{t_1} x^{q_2} : q_i \geq 1, t_j \geq 2\}.$$

Lemma 5.7. *Let $x \in \mathcal{A}$ and let s, s' be sandwiches with $x \notin \text{con}(\bar{s}) = \text{con}(\bar{s}')$. Suppose that $x s x \approx x s' x \in \text{id}(\mathbb{L} \times \mathbb{Z}_n^{\mathbb{R}})$. Then, $s \approx s' \in \text{id}(\mathbb{L} \times \mathbb{Z}_n^{\mathbb{R}})$.*

Proof. This result has been established for the case $\mathbb{R} = 1$ ([10, Lemma 5.8]), but its proof is independent of the value of \mathbb{R} . Hence, the present lemma holds. □

Lemma 5.8. *Suppose that $r \approx r' \in \text{id}_{\text{REF}}(\mathbb{L} \times \mathbb{Z}_n^{\mathbb{R}})$, where*

$$r = x \left(\prod_{i=1}^k p_i \right) x^{\otimes}$$

is the refined sandwich in (5.1). Then,

$$r' = x \left(\prod_{i=1}^k p'_i \right) x^{\otimes}$$

for some $p'_1, p'_2, \dots, p'_k \in (\mathcal{A} \cup \mathcal{A}^*)^+$ such that $x, p'_1, p'_2, \dots, p'_k$ are pairwise disjoint, $\text{con}(\bar{p}_i) = \text{con}(\bar{p}'_i)$ for all i , and each p'_i is either a singleton or a sandwich. Furthermore, for each i ,

- (i) p_i and p'_i are either both singletons or both sandwiches,
- (ii) if p_i and p'_i are both singletons, then $p_i = p'_i$,
- (iii) if p_i and p'_i are both sandwiches, then $p_i \approx p'_i \in \text{id}_{\text{SAN}}(\mathbb{L} \times \mathbb{Z}_n^{\mathbb{R}})$.

Proof. It follows from the assumption and Lemmas 5.1–5.3 that $r' = x u' x^{\otimes}$ for some $u' \in (\mathcal{A} \cup \mathcal{A}^*)^+$ such that

(A) $x \notin \text{con}(\bar{u}') = \text{con}(\bar{p}_1 \cdot \bar{p}_2 \cdots \bar{p}_k)$.

Let $u' = \prod_{i=1}^{k'} p'_i$ be the natural decomposition of u' , so that

$$r' = x \left(\prod_{i=1}^{k'} p'_i \right) x^{\otimes},$$

where

(B) each \mathbf{p}'_i is either a singleton or a sandwich and $x, \mathbf{p}'_1, \mathbf{p}'_2, \dots, \mathbf{p}'_{k'}$ are pairwise disjoint. It is first shown that $k = k'$ and $\text{con}(\overline{\mathbf{p}}_i) = \text{con}(\overline{\mathbf{p}'_i})$ for all i .

Let $\varphi : \mathcal{A} \rightarrow \mathcal{A}^+$ denote the substitution given by

$$z\varphi = \begin{cases} y_i^n & \text{if } z \in \text{con}(\overline{\mathbf{p}}_i), \\ x^n & \text{otherwise.} \end{cases}$$

Note that for any $w \in (\mathcal{A} \cup \mathcal{A}^*)^+$, the image $w\varphi$ belongs to

$$(\mathcal{A}\varphi \cup (\mathcal{A}\varphi)^*)^+ = \{x^n, y_1^n, y_2^n, \dots, y_k^n, (x^n)^*, (y_1^n)^*, (y_2^n)^*, \dots, (y_k^n)^*\}^+.$$

Therefore, identity (3.1d) can be used to convert $w\varphi$ into the plain word $\overline{w\varphi}$ in $\{x^n, y_1^n, y_2^n, \dots, y_k^n\}^+$. Hence,

$$\mathbf{r}\varphi \stackrel{(3.1d)}{\approx} \overline{\mathbf{r}\varphi} = x^n \left(\prod_{i=1}^k y_i^{n|\mathbf{p}_i|} \right) x^n \quad \text{and} \quad \mathbf{r}'\varphi \stackrel{(3.1d)}{\approx} \overline{\mathbf{r}'\varphi} = x^n \left(\prod_{i=1}^{k'} \overline{\mathbf{p}'_i\varphi} \right) x^n.$$

Since $\overline{\mathbf{r}\varphi} \approx \overline{\mathbf{r}'\varphi} \in \text{id}(\mathbf{L})$ and $\overline{\mathbf{r}\varphi} \in \mathfrak{P}_k^\uparrow$, Lemma 5.6 implies that $\overline{\mathbf{r}'\varphi} \in \mathfrak{P}_k^\uparrow \cup \mathfrak{P}_k^\downarrow$, that is,

(C) $x^n \left(\prod_{i=1}^{k'} \overline{\mathbf{p}'_i\varphi} \right) x^n \in \mathfrak{P}_k^\uparrow \cup \mathfrak{P}_k^\downarrow$.

Furthermore, since

$$\text{con}(\overline{\mathbf{p}'_i\varphi}) \subseteq \text{con}(\overline{\mathbf{u}'\varphi}) \stackrel{(A)}{=} \text{con}(\overline{(\mathbf{p}_1\mathbf{p}_2 \cdots \mathbf{p}_k)\varphi}) = \{y_1, y_2, \dots, y_k\},$$

it follows that

(D) $\text{con}(\overline{\mathbf{p}'_i\varphi}) \subseteq \{y_1, y_2, \dots, y_k\}$ for each $i \in \{1, 2, \dots, k'\}$.

Recall that each \mathbf{p}'_i is either a singleton or a sandwich. If \mathbf{p}'_i is a singleton, then clearly $\text{con}(\overline{\mathbf{p}'_i\varphi}) = \{y_j\}$ for some j . Suppose that \mathbf{p}'_i is a sandwich. Then, \mathbf{p}'_i is connected, so that $\overline{\mathbf{p}'_i\varphi}$ is a connected factor of $\overline{\mathbf{r}'\varphi} \in \mathfrak{P}_k^\uparrow \cup \mathfrak{P}_k^\downarrow$. The connected factors of words in $\mathfrak{P}_k^\uparrow \cup \mathfrak{P}_k^\downarrow$ are

$$x^{q_1} y_1^{t_1} y_2^{t_2} \cdots y_k^{t_k} x^{q_2}, \quad x^{q_1} y_k^{t_k} y_{k-1}^{t_{k-1}} \cdots y_1^{t_1} x^{q_2}, \quad q_i \geq 1, t_j \geq 2,$$

and

$$x^t, y_1^t, y_2^t, \dots, y_k^t, \quad t \geq 2.$$

But since $x \notin \text{con}(\overline{\mathbf{p}'_i\varphi})$ by (D), the word $\overline{\mathbf{p}'_i\varphi}$ can only be one of $y_1^t, y_2^t, \dots, y_k^t$, so that $\text{con}(\overline{\mathbf{p}'_i\varphi}) = \{y_j\}$ for some j . Hence, regardless of whether \mathbf{p}'_i is a singleton or a sandwich,

(E) $\text{con}(\overline{\mathbf{p}'_i\varphi}) = \{y_j\}$ for some j .

It follows that $\text{con}(\overline{\mathbf{p}'_i}) \subseteq \text{con}(\overline{\mathbf{p}}_j)$ for some j . By a symmetric argument, the inclusion $\text{con}(\overline{\mathbf{p}}_j) \subseteq \text{con}(\overline{\mathbf{p}'_m})$ holds for some m , so that $\text{con}(\overline{\mathbf{p}'_i}) \subseteq \text{con}(\overline{\mathbf{p}}_j) \subseteq \text{con}(\overline{\mathbf{p}'_m})$. Since the words \mathbf{p}'_i and \mathbf{p}'_m are either equal or disjoint, $\mathbf{p}'_i = \mathbf{p}'_m$ is the only possibility, whence $\text{con}(\overline{\mathbf{p}'_i}) = \text{con}(\overline{\mathbf{p}}_j)$. It has just been shown that for each $i \in \{1, 2, \dots, k'\}$, there exists some $j \in \{1, 2, \dots, k\}$ such that $\text{con}(\overline{\mathbf{p}'_i}) = \text{con}(\overline{\mathbf{p}}_j)$. Since $\text{con}(\overline{\mathbf{p}}_1 \cdots \overline{\mathbf{p}}_k) = \text{con}(\overline{\mathbf{p}'_1} \cdots \overline{\mathbf{p}'_{k'}})$ by (A), it follows that

(F) $k = k'$ and

(G) there exists a one-to-one correspondence between

$$\text{con}(\overline{\mathbf{p}}_1), \text{con}(\overline{\mathbf{p}}_2), \dots, \text{con}(\overline{\mathbf{p}}_k) \quad \text{and} \quad \text{con}(\overline{\mathbf{p}'_1}), \text{con}(\overline{\mathbf{p}'_2}), \dots, \text{con}(\overline{\mathbf{p}'_{k'}}).$$

Furthermore, (C) and (F) imply that either

$$x^n \left(\prod_{i=1}^k \overline{\mathbf{p}'_i\varphi} \right) x^n \in \mathfrak{P}_k^\uparrow \quad \text{or} \quad x^n \left(\prod_{i=1}^{k'} \overline{\mathbf{p}'_i\varphi} \right) x^n \in \mathfrak{P}_k^\downarrow.$$

It thus follows from (E) that either

(H) $(\text{con}(\overline{\mathbf{p}'_1}), \text{con}(\overline{\mathbf{p}'_2}), \dots, \text{con}(\overline{\mathbf{p}'_{k'}})) = (\text{con}(\overline{\mathbf{p}}_1), \text{con}(\overline{\mathbf{p}}_2), \dots, \text{con}(\overline{\mathbf{p}}_k))$ or

(H') $(\text{con}(\overline{\mathbf{p}'_1}), \text{con}(\overline{\mathbf{p}'_2}), \dots, \text{con}(\overline{\mathbf{p}'_{k'}})) = (\text{con}(\overline{\mathbf{p}}_k), \text{con}(\overline{\mathbf{p}}_{k-1}), \dots, \text{con}(\overline{\mathbf{p}}_1))$.

Now Lemma 2.1 implies that

(I) $\text{sim}(\overline{\mathbf{r}}) = \text{sim}(\overline{\mathbf{r}'})$ and $\text{non}(\overline{\mathbf{r}}) = \text{non}(\overline{\mathbf{r}'})$.

If $k = 1$, then (H) clearly holds, so assume that $k \geq 2$.

Case 1: The words p_1, p_2, \dots, p_k are all sandwiches. Then, in view of (B) and (I), either (H) or (H') implies that p'_1, p'_2, \dots, p'_k are also sandwiches. Hence, (H) must hold by (R.2).

Case 2: The word p_i is a singleton for some i . Then, $p_i \in \{y, y^*\}$ for some $y \in \text{sim}(\bar{r})$. It follows from either (H) or (H') that $\text{con}(p'_j) = \{y\}$ for some j , whence $p'_j \in \{y, y^*\}$ by (I). If $p_i \neq p'_j$, so that $(p_i, p'_j) \in \{(y, y^*), (y^*, y)\}$, then since the group Z_n^R satisfies $r \approx r'$ and has a unit element, it also satisfies the identity $y^* \approx y$; this is impossible because $g^* = g^R \neq g$. Therefore, $p_i = p'_j \in \{y, y^*\}$. Now since $k \geq 2$, either $1 < i$ or $i < k$. By symmetry, assume that $1 < i$. Let $\chi : \mathcal{A} \rightarrow L$ denote the substitution given by

$$z\chi = \begin{cases} e & \text{if } z \in \text{con}(\overline{p_1} \cdot \overline{p_2} \cdots \overline{p_{i-1}}), \\ ef & \text{if } z = y \text{ and } p_i = y, \\ fe & \text{if } z = y \text{ and } p_i = y^*, \\ f & \text{otherwise.} \end{cases}$$

Then,

$$\begin{aligned} r\chi &= (x \cdot p_1 \cdot p_2 \cdots p_{i-1} \cdot p_i \cdot p_{i+1} \cdots p_k \cdot x)\chi \\ &= f \cdot e \cdot e \cdots e \cdot ef \cdot f \cdots f \cdot f \\ &= fef. \end{aligned} \tag{5.2}$$

If (H') holds, then $r'\chi$ is the product (5.2) in reverse order, that is,

$$r'\chi = f \cdot f \cdots f \cdot ef \cdot e \cdot e \cdots e \cdot f = 0.$$

But this is impossible, so that (H') cannot hold. Therefore, (H) must hold.

Hence, (H) holds in any case. It then follows from (B) and (I) that (i) and (ii) hold. It remains to verify that (iii) also holds. Suppose that p_i and p'_i are sandwiches. Let $\psi : \mathcal{A} \rightarrow \mathcal{A}$ denote the substitution given by

$$z\psi = \begin{cases} z & \text{if } z \in \text{con}(\overline{p_i}) = \text{con}(\overline{p'_i}), \\ x^n & \text{otherwise.} \end{cases}$$

Then, the deductions (3.1a) $\vdash \{x(r\psi)x \approx xp_i x, x(r'\psi)x \approx xp'_i x\}$ hold, so that the deduction $\{(3.1a), r \approx r'\} \vdash xp_i x \approx xp'_i x$ also holds. It now follows from Lemma 3.2 that $xp_i x \approx xp'_i x \in \text{id}(L \times Z_n^R)$. Hence, $p_i \approx p'_i \in \text{id}(L \times Z_n^R)$ by Lemma 5.7. \square

6 Proof of Theorem 3.1

In this section, the following statement is established for each $m \geq 1$.

(\mathcal{S}_m) If $s \approx s' \in \text{id}_{\text{SAN}}(L \times Z_n^R)$ with $|\text{non}(\bar{s})| \leq m$, then (3.1) $\vdash s \approx s'$.

Hence, the deduction (3.1) $\vdash \text{id}_{\text{SAN}}(L \times Z_n^R)$ holds. It follows from Lemma 4.4 that the deduction

$$\{(3.1)\} \cup \text{id}_{\text{SAN}}(L \times Z_n^R) \vdash \text{id}(L \times Z_n^R)$$

holds. Therefore, the proof of Theorem 3.1 is complete.

Lemma 6.1. *The statement (\mathcal{S}_1) holds.*

Proof. Suppose that $s \approx s' \in \text{id}_{\text{SAN}}(L \times Z_n^R)$ with $|\text{non}(\bar{s})| = 1$. Then, by Lemma 5.5, there exists some finite $\Sigma \subseteq \text{id}_{\text{REF}}(L \times Z_n^R)$ such that

(A) $\{(3.1), s \approx s'\} \sim \{(3.1)\} \cup \Sigma$ and

(B) each $r \approx r' \in \Sigma$ satisfies $|\text{sim}(\bar{r})| \leq |\text{sim}(\bar{s})|$ and $|\text{non}(\bar{r})| \leq |\text{non}(\bar{s})| = 1$.

Consider any $r \approx r' \in \Sigma$. Generality is not lost by assuming that $r = x(\prod_{i=1}^k p_i)x^\otimes$ is the refined sandwich in (5.1). Then, Lemma 5.8 implies that $r' = x(\prod_{i=1}^k p'_i)x^\otimes$ for some $p'_i \in (\mathcal{A} \cup \mathcal{A}^*)^+$ such that $\text{con}(\overline{p_i}) = \text{con}(\overline{p'_i})$

for all i . Since $1 = |\text{non}(\bar{r})| = |\text{non}(\bar{r}')|$ by (B) and Lemma 2.1, it follows from (i) and (ii) of Lemma 5.8 that \mathbf{p}_i and \mathbf{p}'_i are the same singleton. Hence, the identity $\mathbf{r} \approx \mathbf{r}'$ is trivial.

Since the identity $\mathbf{r} \approx \mathbf{r}'$ is arbitrary in Σ , every identity in Σ is trivial. The deduction (3.1) $\vdash \mathbf{s} \approx \mathbf{s}'$ thus follows from (A). □

Lemma 6.2. *Suppose that the statement (\mathcal{S}_m) holds. Then, the statement (\mathcal{S}_{m+1}) also holds.*

Proof. Suppose that $\mathbf{s} \approx \mathbf{s}' \in \text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ with $|\text{non}(\mathbf{s})| = m + 1$. Then, by Lemma 5.5, there exists some finite $\Sigma \subseteq \text{id}_{\text{REF}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ such that

(A) $\{(3.1), \mathbf{s} \approx \mathbf{s}'\} \sim \{(3.1)\} \cup \Sigma$ and

(B) each $\mathbf{r} \approx \mathbf{r}' \in \Sigma$ satisfies $|\text{sim}(\bar{\mathbf{r}})| \leq |\text{sim}(\bar{\mathbf{s}})|$ and $|\text{non}(\bar{\mathbf{r}})| \leq |\text{non}(\bar{\mathbf{s}})| = m + 1$.

Consider any $\mathbf{r} \approx \mathbf{r}' \in \Sigma$. Generality is not lost by assuming that $\mathbf{r} = x(\prod_{i=1}^k \mathbf{p}_i)x^{\otimes}$ is the refined sandwich in (5.1). Then, Lemma 5.8 implies that $\mathbf{r}' = x(\prod_{i=1}^k \mathbf{p}'_i)x^{\otimes}$ for some $\mathbf{p}'_i \in (\mathcal{A} \cup \mathcal{A}^*)^+$ such that $\text{con}(\bar{\mathbf{p}}_i) = \text{con}(\bar{\mathbf{p}}'_i)$ for all i . By (i) of Lemma 5.8, the words \mathbf{p}_i and \mathbf{p}'_i are both singletons or both sandwiches.

Case 1: The words \mathbf{p}_i and \mathbf{p}'_i are singletons. Then, one has $\mathbf{p}_i = \mathbf{p}'_i$ by (ii) of Lemma 5.8, so that the deduction (3.1) $\vdash \mathbf{p}_i \approx \mathbf{p}'_i$ holds vacuously.

Case 2: The words \mathbf{p}_i and \mathbf{p}'_i are both sandwiches. Then, $\mathbf{p}_i \approx \mathbf{p}'_i \in \text{id}_{\text{SAN}}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ by (iii) of Lemma 5.8. Since $|\text{non}(\bar{\mathbf{p}}_i)| < |\text{non}(\bar{\mathbf{r}})| \leq m + 1$ by (B), the deduction (3.1) $\vdash \mathbf{p}_i \approx \mathbf{p}'_i$ follows from (\mathcal{S}_m) .

Therefore, the deduction (3.1) $\vdash \mathbf{p}_i \approx \mathbf{p}'_i$ holds in any case. Since

$$\mathbf{r} = x\left(\prod_{i=1}^k \mathbf{p}_i\right)x^{\otimes} \stackrel{(3.1)}{\approx} x\left(\prod_{i=1}^k \mathbf{p}'_i\right)x^{\otimes} = \mathbf{r}'$$

the deduction (3.1) $\vdash \mathbf{r} \approx \mathbf{r}'$ also holds. The identity $\mathbf{r} \approx \mathbf{r}'$ is arbitrary in Σ , so that (3.1) $\vdash \Sigma$. Consequently, the deduction (3.1) $\vdash \mathbf{s} \approx \mathbf{s}'$ follows from (A). □

7 Irredundant basis property

7.1 Terms, identities, and deducibility

The set $T_{\mathcal{A}}$ of terms over \mathcal{A} is the smallest set that satisfies all of the following:

- $\mathcal{A} \subseteq T_{\mathcal{A}}$,
- if $\mathbf{t}_1, \mathbf{t}_2 \in T_{\mathcal{A}}$, then $\mathbf{t}_1\mathbf{t}_2 \in T_{\mathcal{A}}$,
- if $\mathbf{t} \in T_{\mathcal{A}}$, then $\mathbf{t}^* \in T_{\mathcal{A}}$.

The subterms of a term \mathbf{t} are defined as follows:

- \mathbf{t} is a subterm of \mathbf{t} ,
- if $\mathbf{t}_1\mathbf{t}_2$ is a subterm of \mathbf{t} , where $\mathbf{t}_1, \mathbf{t}_2 \in T_{\mathcal{A}}$, then \mathbf{t}_1 and \mathbf{t}_2 are subterms of \mathbf{t} ,
- if \mathbf{u}^* is a subterm of \mathbf{t} , where $\mathbf{u} \in T_{\mathcal{A}}$, then \mathbf{u} is a subterm of \mathbf{t} .

Remark 7.1. Note the following.

- (i) The inclusion $(\mathcal{A} \cup \mathcal{A}^*)^+ \subset T_{\mathcal{A}}$ holds.
- (ii) The identities (inv) can be used to convert any term $\mathbf{t} \in T_{\mathcal{A}}$ into a unique word in $(\mathcal{A} \cup \mathcal{A}^*)^+$; denote this unique word by $[\mathbf{t}]$.
- (iii) If \mathbf{u} is a subterm of a term $\mathbf{t} \in T_{\mathcal{A}}$, then either $[\mathbf{u}]$ or $[\mathbf{u}^*]$ is a factor of the word $[\mathbf{t}]$.

For the remainder of this article, identities $\mathbf{t} \approx \mathbf{t}'$ are formed by terms $\mathbf{t}, \mathbf{t}' \in T_{\mathcal{A}}$. An identity $\mathbf{w} \approx \mathbf{w}'$ formed by words $\mathbf{w}, \mathbf{w}' \in (\mathcal{A} \cup \mathcal{A}^*)^+$ is called a *word identity*. The objective of Sections 3–6 was to prove that identities (3.1) constitute a basis for the involution semigroup $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$. In view of (ii) of Remark 7.1, this task was achievable by working mainly with word identities. However, the situation is different in the present section, where the goal is to

(‡) extract from (3.1) an infinite irredundant basis for $\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}}$;

see Theorem 7.4. Each identity from this extracted basis is shown to be undeducible from all other identities in the basis, and this involves examining deduction sequences that generally contain terms instead of only words. Working with only word identities is thus insufficient.

Due to (‡), the concept of deducibility of identities, first defined in Section 2, has to be treated more formally. An identity $x \approx y$ is *directly deducible* from an identity $u \approx v$ if there exists some substitution $\varphi : \mathcal{A} \rightarrow T_{\mathcal{A}}$ such that $u\varphi$ is a subterm of x , and replacing this subterm of x with $v\varphi$ results in the term y . By Birkhoff’s completeness theorem of equational logic [2], an identity $x \approx y$ is *deducible* from a set Σ of identities if there exists a sequence

$$x = t_1, t_2, \dots, t_r = y$$

of terms, where each identity $t_i \approx t_{i+1}$ is directly deducible from some identity in Σ .

Recall that the number of times a letter $x \in \mathcal{A}$ occurs in a word $w \in (\mathcal{A} \cup \mathcal{A}^*)^+$ is denoted by $\text{occ}(x, \bar{w})$. The number of times $x \in \mathcal{A}$ occurs in a term $t \in T_{\mathcal{A}}$ is $\text{occ}(x, \overline{[t]})$. For instance, if $t = x(y^*x^2)^*(zx^*y)^* \in T_{\mathcal{A}}$, then

$$\overline{[t]} = \overline{x(x^*)^2yy^*xz^*} = x^3y^2xz,$$

so that $\text{occ}(x, \overline{[t]}) = 4$, $\text{occ}(y, \overline{[t]}) = 2$, and $\text{occ}(z, \overline{[t]}) = 1$. The following result is easily seen to hold.

Lemma 7.2. *Suppose that $t \approx t'$ is any identity deducible from (inv). Then, one has $[t] = [t']$.*

7.2 Identities (3.1w)

For each $m \geq 2$, let

$$\mathbb{B}_m = \{(b_1, b_2, \dots, b_m) : b_1, b_2, \dots, b_m \in \{0, 1\}\}$$

denote the set of all binary vectors of dimension m . The vectors in \mathbb{B}_m are lexicographically ordered by $<$ as

$$(b_1, b_2, \dots, b_m) < (b'_1, b'_2, \dots, b'_m)$$

if there exists a least $\ell \in \{1, 2, \dots, m\}$ such that $b_\ell < b'_\ell$ and $b_i = b'_i$ for any $i < \ell$. If $v = (b_1, b_2, \dots, b_m) \in \mathbb{B}_m$, then the *dual* of v is $\delta v = (b_m, b_{m-1}, \dots, b_1)$. If $\delta v = v$, then v is a *palindrome*. The set \mathbb{B}_m can be partitioned into the three subsets

$$\mathbb{B}_m^= = \{v \in \mathbb{B}_m : v = \delta v\}, \quad \mathbb{B}_m^< = \{v \in \mathbb{B}_m : v < \delta v\}, \quad \mathbb{B}_m^> = \{v \in \mathbb{B}_m : v > \delta v\}.$$

Each vector $v = (b_1, b_2, \dots, b_m) \in \mathbb{B}_m$ is associated with the words

$$v^\uparrow = x \left(\prod_{i=1}^m (y_i h_i^{b_i} y_i^*) \right) x^* = x \cdot y_1 h_1^{b_1} y_1^* \cdot y_2 h_2^{b_2} y_2^* \cdots y_m h_m^{b_m} y_m^* \cdot x^*,$$

$$v^\downarrow = x \left(\prod_{i=m}^1 (y_i h_i^{b_i} y_i^*) \right) x^* = x \cdot y_m h_m^{b_m} y_m^* \cdot y_{m-1} h_{m-1}^{b_{m-1}} y_{m-1}^* \cdots y_1 h_1^{b_1} y_1^* \cdot x^*,$$

where $h_i^0 = \emptyset$ and $h_i^1 = h_i$. Note that

$$\bigcup_{m \geq 2} \{v^\uparrow \approx v^\downarrow : v \in \mathbb{B}_m\} = \{(3.1w)\}.$$

In Section 7.3, some identities from (3.1w) are chosen to form an irredundant basis for the involution semigroup $L \times Z_n^R$. But for each $m \geq 2$, the identities in $\{v^\uparrow \approx v^\downarrow : v \in \mathbb{B}_m\}$ are not irredundant; if v is not a palindrome, then v and δv are distinct vectors such that the associated identities $v^\uparrow \approx v^\downarrow$ and $(\delta v)^\uparrow \approx (\delta v)^\downarrow$ are equivalent. This redundancy can be eliminated by choosing identities $v^\uparrow \approx v^\downarrow$ with v taken from only $\mathbb{B}_m^=$ or $\mathbb{B}_m^<$. Hence, the equivalence

$$(3.1w) \sim \bigcup_{m \geq 2} \{v^\uparrow \approx v^\downarrow : v \in \mathbb{B}_m^= \cup \mathbb{B}_m^<\}$$

holds, where no two identities on the right are equivalent.

Lemma 7.3. Let $v = (b_1, b_2, \dots, b_m) \in \mathbb{B}_m$. Suppose that $v^\dagger \approx w \in \text{id}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$ for some word $w \in (\mathcal{A} \cup \mathcal{A}^*)^+$. Then, either $w = v^\dagger$ or $w = v^\downarrow$.

Proof. By Lemma 2.1,

(A) $\text{sim}(\bar{w}) = \{h_1^{b_1}, h_2^{b_2}, \dots, h_m^{b_m}\}$ and $\text{non}(\bar{w}) = \{x, y_1, y_2, \dots, y_m\}$.

Let $\varphi : \mathcal{A} \rightarrow L$ denote the substitution given by

$$z\varphi = \begin{cases} fe & \text{if } z = x, \\ e & \text{otherwise.} \end{cases}$$

Then, $v^\dagger\varphi = fef = w\varphi$, so that $w\varphi$ is a product of the form $fe \cdot e \cdot e \cdots e \cdot ef$, whence

(B) $w = x\mathbf{u}x^*$ for some $\mathbf{u} \in (\mathcal{A} \cup \mathcal{A}^*)^+$ such that $x \notin \text{con}(\bar{\mathbf{u}})$.

Consider any fixed $i \in \{1, 2, \dots, m\}$. Let $\chi : \mathcal{A} \rightarrow L$ denote the substitution given by

$$z\chi = \begin{cases} fe & \text{if } z = y_i, \\ e & \text{if } z = h_i, \\ f & \text{otherwise.} \end{cases}$$

Then, $v^\dagger\chi = fef = w\chi$.

There are two cases to consider.

Case 1. If $b_i = 1$, then $h_i \in \text{sim}(\bar{w})$ by (A), so that

$$w\chi = f \cdot f \cdots f \cdot fe \cdot e \cdot ef \cdot f \cdot f \cdots f.$$

Hence, (A) and (B) imply that $w = x\mathbf{u}_1 y_i h_i^{\otimes} y_i^* \mathbf{u}_2 x^*$ for some $\otimes \in \{1, *\}$ and $\mathbf{u}_1, \mathbf{u}_2 \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $x, y_i, h_i \notin \text{con}(\bar{\mathbf{u}}_1 \cdot \bar{\mathbf{u}}_2)$. Since $v^\dagger \approx w \in \text{id}(\mathbf{Z}_n^{\mathbb{R}})$ and $\mathbf{Z}_n^{\mathbb{R}}$ has a unit element, it follows that $h_i \approx h_i^{\otimes} \in \text{id}(\mathbf{Z}_n^{\mathbb{R}})$. But $g \neq g^*$ in $\mathbf{Z}_n^{\mathbb{R}}$, so that $\otimes = 1$. Thus $w = x\mathbf{u}_1 y_i h_i^1 y_i^* \mathbf{u}_2 x^* = x\mathbf{u}_1 y_i h_i^b y_i^* \mathbf{u}_2 x^*$.

Case 2. If $b_i = 0$, then $h_i \notin \text{con}(\bar{w})$ by (A), so that

$$w\chi = f \cdot f \cdots f \cdot fe \cdot ef \cdot f \cdot f \cdots f.$$

Hence, (A) and (B) imply that $w = x\mathbf{u}_1 y_i y_i^* \mathbf{u}_2 x^* = x\mathbf{u}_1 y_i h_i^b y_i^* \mathbf{u}_2 x^*$ for some $\mathbf{u}_1, \mathbf{u}_2 \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $x, y_i \notin \text{con}(\bar{\mathbf{u}}_1 \cdot \bar{\mathbf{u}}_2)$.

Therefore, in any case, $w = x\mathbf{u}_1 y_i h_i^{b_i} y_i^* \mathbf{u}_2 x^*$ for some $\mathbf{u}_1, \mathbf{u}_2 \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ such that $x, y_i, h_i \notin \text{con}(\bar{\mathbf{u}}_1 \cdot \bar{\mathbf{u}}_2)$. Since i is arbitrary, it follows that $w = x\mathbf{v}x^*$, where \mathbf{v} is a product of $y_1 h_1^{b_1} y_1^*, y_2 h_2^{b_2} y_2^*, \dots, y_m h_m^{b_m} y_m^*$ in some order. Let π denote the permutation on $\{1, 2, \dots, m\}$ such that

$$w = x \left(\prod_{i=1}^m (y_{i\pi} h_{i\pi}^{b_{i\pi}} y_{i\pi}^*) \right) x^*.$$

Let $\psi : \mathcal{A} \rightarrow \mathcal{A}^+$ denote the substitution given by

$$z\psi = \begin{cases} y_i^n & \text{if } z \in \{y_i, h_i\}, \\ x^n & \text{otherwise.} \end{cases}$$

Then,

$$v^\dagger\psi \stackrel{(3.1a), (3.1d)}{\approx} x^n \left(\prod_{i=1}^m y_i^n \right) x^n \in \mathfrak{P}_m^\dagger \quad \text{and} \quad w\psi \stackrel{(3.1a), (3.1d)}{\approx} x^n \left(\prod_{i=1}^m y_{i\pi}^n \right) x^n.$$

Since $x^n \left(\prod_{i=1}^m y_i^n \right) x^n \approx x^n \left(\prod_{i=1}^m y_{i\pi}^n \right) x^n \in \text{id}(\mathbf{L} \times \mathbf{Z}_n^{\mathbb{R}})$, it follows from Lemma 5.6 that $x^n \left(\prod_{i=1}^m y_{i\pi}^n \right) x^n \in \mathfrak{P}_m^\dagger \cup \mathfrak{P}_m^\downarrow$, whence

$$(1\pi, 2\pi, \dots, m\pi) \in \{(1, 2, \dots, m), (m, m-1, \dots, 1)\}.$$

Consequently, either $w = v^\dagger$ or $w = v^\downarrow$. □

7.3 An irredundant basis from (3.1)

Theorem 7.4. *An infinite irredundant basis for the involution semigroup $L \times Z_n^R$ can be formed from the identities in (3.1).*

Recall from Section 7.2 that identities (3.1w) and

$$\Omega = \bigcup_{m \geq 2} \{v^\uparrow \approx v^\downarrow : v \in B_m^\bar{} \cup B_m^<\}$$

are equivalent. Since the set $\Theta = \{(3.1a), (3.1b), \dots, (3.1v)\}$ is finite, it follows from Theorem 3.1 that Θ contains some minimal subset Θ_{\min} such that $\Theta_{\min} \cup \Omega$ is a basis for the involution semigroup $L \times Z_n^R$. Let

$$v_{\text{fix}} = (b_1, b_2, \dots, b_m) \in B_m^\bar{} \cup B_m^<$$

be a fixed vector. In the remainder of this subsection, it is shown that the identity

$$v_{\text{fix}}^\uparrow \approx v_{\text{fix}}^\downarrow$$

in Ω is not deducible from

$$\{(\text{inv})\} \cup \Theta_{\min} \cup (\Omega \setminus \{v_{\text{fix}}^\uparrow \approx v_{\text{fix}}^\downarrow\}). \tag{7.1}$$

Since Ω is infinite, the set $\Theta_{\min} \cup \Omega$ is an infinite irredundant basis for the involution semigroup $L \times Z_n^R$. This completes the proof of Theorem 7.4.

Seeking a contradiction, suppose that the identity $v_{\text{fix}}^\uparrow \approx v_{\text{fix}}^\downarrow$ is deducible from identities (7.1). Then, there exists a sequence

$$v_{\text{fix}}^\uparrow = t_1, t_2, \dots, t_r = v_{\text{fix}}^\downarrow$$

of terms, where each identity $t_i \approx t_{i+1}$ is directly deducible from some identity $u_i \approx v_i$ in (7.1). If every identity $u_i \approx v_i$ is from (inv), then it follows from Lemma 7.2 that $v_{\text{fix}}^\uparrow = [t_i]$ for all i , whence the contradiction $v_{\text{fix}}^\uparrow = [t_r] = v_{\text{fix}}^\downarrow$ is obtained. Therefore, some $u_i \approx v_i$ is not from (inv); let $\ell \geq 1$ be the least possible index such that $u_\ell \approx v_\ell$ is not from (inv). Then,

(I) $u_\ell \approx v_\ell$ is from $\Theta_{\min} \cup (\Omega \setminus \{v_{\text{fix}}^\uparrow \approx v_{\text{fix}}^\downarrow\})$

while $u_1 \approx v_1, u_2 \approx v_2, \dots, u_{\ell-1} \approx v_{\ell-1}$ are from (inv), whence

(II) $v_{\text{fix}}^\uparrow \approx t_\ell$ is deducible from (inv).

In what follows, it is shown that the identity $u_\ell \approx v_\ell$ belongs to neither Θ_{\min} nor $\Omega \setminus \{v_{\text{fix}}^\uparrow \approx v_{\text{fix}}^\downarrow\}$; this and (I) imply the required contradiction.

Lemma 7.5. *The following hold.*

(i) $[t_\ell] = v_{\text{fix}}^\uparrow$.

(ii) *The word $[t_\ell]$ does not contain any of the factors*

$$x^{\otimes 1} h_1 x^{\otimes 1}, \tag{7.2a}$$

$$x^{\otimes 1} h_1 x^{\otimes 2} h_2 x^{\otimes 3}, \tag{7.2b}$$

$$x^{\otimes 1} h_1 y^{\otimes 2} h_2 x^{\otimes 3} h_3 y^{\otimes 4}, \tag{7.2c}$$

$$x^{\otimes 1} h_1 y^{\otimes 2} h_2 z^{\otimes 3} h_3 z^{\otimes 4} h_4 y^{\otimes 5} h_5 x^{\otimes 6}, \tag{7.2d}$$

$$x^{\otimes 1} h_1 y^{\otimes 2} h_2 z^{\otimes 3} t^{\otimes 4} h_3 y^{\otimes 5} h_4 x^{\otimes 6}, \tag{7.2e}$$

where $x, y, z, t \in \mathcal{A}$, $h_i \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$, and $\otimes_i \in \{1, *\}$.

[Correction added after online publication 11 September 2015: For the readers convenience equation (7.2a) has been changed from $x^{\otimes 1} h_1 x^{\otimes 2}$ to $x^{\otimes 1} h_1 x^{\otimes 1}$.]

Proof. Part (i) follows from (II) and the observation made in (ii) of Remark 7.1, while part (ii) is a consequence of part (i). □

Since the identity $t_\ell \approx t_{\ell+1}$ is directly deducible from $u_\ell \approx v_\ell$, there exists some substitution $\varphi : \mathcal{A} \rightarrow T_{\mathcal{A}}$ such that $u_\ell \varphi$ is a subterm of t_ℓ .

Lemma 7.6. *The identity $u_\ell \approx v_\ell$ cannot be from Θ_{\min} .*

Proof. Suppose that $u_\ell \approx v_\ell$ is an identity from (3.1i)–(3.1v). Then,

$$u_\ell = h_0 x^{\otimes_1} h_1 y^{\otimes_2} h_2 x^{\otimes_3} h_3 y^{\otimes_4} h_4$$

for some $h_i \in (\mathcal{A} \cup \mathcal{A}^*)^+ \cup \{\emptyset\}$ and $\otimes_i \in \{1, *\}$. Let $z \in \text{con}(\overline{[x\varphi]})$ and $t \in \text{con}(\overline{[y\varphi]})$, so that

$$[u_\ell \varphi] = [h_0 \varphi] \underbrace{(\dots z^{\otimes_1} \dots)}_{[(x\varphi)^{\otimes_1}]} [h_1 \varphi] \underbrace{(\dots t^{\otimes_2} \dots)}_{[(y\varphi)^{\otimes_2}]} [h_2 \varphi] \underbrace{(\dots z^{\otimes_3} \dots)}_{[(x\varphi)^{\otimes_3}]} [h_3 \varphi] \underbrace{(\dots t^{\otimes_4} \dots)}_{[(y\varphi)^{\otimes_4}]} [h_4 \varphi]$$

for some $\otimes'_i \in \{1, *\}$. Since $u_\ell \varphi$ is a subterm of t_ℓ , as observed in (iii) of Remark 7.1, either $[u_\ell \varphi]$ or $[(u_\ell \varphi)^*]$ is a factor of the word $[t_\ell]$, whence $[t_\ell]$ contains a factor of the form (7.2c). But this is impossible by (ii) of Lemma 7.5.

If $u_\ell \approx v_\ell$ is an identity from (3.1a)–(3.1h), then a similar argument shows that the word $[t_\ell]$ contains a factor of the form (7.2a) or (7.2b). This is again impossible by (ii) of Lemma 7.5. \square

Lemma 7.7. *The identity $u_\ell \approx v_\ell$ cannot be from $\Omega \setminus \{V_{\text{fix}}^\uparrow \approx V_{\text{fix}}^\downarrow\}$.*

Proof. It suffices to assume that the identity $u_\ell \approx v_\ell$ belongs to Ω and then show that it is precisely $V_{\text{fix}}^\uparrow \approx V_{\text{fix}}^\downarrow$. Suppose that $u_\ell \approx v_\ell$ is the identity $W^\uparrow \approx W^\downarrow$ in Ω , where $W = (c_1, c_2, \dots, c_n) \in \mathbb{B}_n^\equiv \cup \mathbb{B}_n^<$ for some $n \geq 2$. Then, either $u_\ell \varphi = W^\uparrow \varphi$ or $u_\ell \varphi = W^\downarrow \varphi$ is a subterm of t_ℓ ; by symmetry, it suffices to assume that

$$u_\ell \varphi = W^\uparrow \varphi = (x\varphi) \left(\prod_{i=1}^n ((y_i \varphi)(h_i^{c_i} \varphi)(y_i \varphi)^*) \right) (x\varphi)^*$$

is a subterm of t_ℓ .

Suppose that the word $[x\varphi]$ is non-singleton. Then, $\overline{[x\varphi]} = \dots x_1 x_2 \dots$ for some $x_1, x_2 \in \mathcal{A}$. Choose any $q \in \text{con}(\overline{[y_1 \varphi]})$. Then,

$$\begin{aligned} [W^\uparrow \varphi] &= [x\varphi][y_1 \varphi][h_1^{c_1} \varphi][(y_1 \varphi)^*] \dots [(x\varphi)^*] \\ &= \underbrace{(\dots x_1^{\otimes_1} x_2^{\otimes_2} \dots)}_{[x\varphi]} \underbrace{(\dots q^{\otimes_3} \dots)}_{[y_1 \varphi]} [h_1^{c_1} \varphi] \underbrace{(\dots q^{\otimes_4} \dots)}_{[(y_1 \varphi)^*]} \dots \underbrace{(\dots x_2^{\otimes_5} x_1^{\otimes_6} \dots)}_{[(x\varphi)^*]} \end{aligned}$$

for some $\otimes_i \in \{1, *\}$. Since $W^\uparrow \varphi$ is a subterm of t_ℓ , as observed in (iii) of Remark 7.1, either $[W^\uparrow \varphi]$ or $[(W^\uparrow \varphi)^*]$ is a factor of the word $[t_\ell]$, whence $[t_\ell]$ contains a factor of the form (7.2b) or (7.2d), depending on whether or not the letters x_1, x_2 , and q are distinct. But this is impossible by (ii) of Lemma 7.5. Therefore,

(A) the word $[x\varphi]$ is a singleton.

Suppose that the word $[y_i \varphi]$ is non-singleton. Then, $\overline{[y_i \varphi]} = \dots x_1 x_2 \dots$ for some $x_1, x_2 \in \mathcal{A}$. Choose any $q \in \text{con}(\overline{[x\varphi]})$. Then,

$$\begin{aligned} [W^\uparrow \varphi] &= [x\varphi] \dots [y_i \varphi][h_i^{c_i} \varphi][(y_i \varphi)^*] \dots [(x\varphi)^*] \\ &= \underbrace{(\dots q^{\otimes_1} \dots)}_{[x\varphi]} \dots \underbrace{(\dots x_1^{\otimes_2} x_2^{\otimes_3} \dots)}_{[y_i \varphi]} [h_i^{c_i} \varphi] \underbrace{(\dots x_2^{\otimes_4} x_1^{\otimes_5} \dots)}_{[(y_i \varphi)^*]} \dots \underbrace{(\dots q^{\otimes_6} \dots)}_{[(x\varphi)^*]} \end{aligned}$$

for some $\otimes_j \in \{1, *\}$. Since $W^\uparrow \varphi$ is a subterm of t_ℓ , by (iii) of Remark 7.1, either $[W^\uparrow \varphi]$ or $[(W^\uparrow \varphi)^*]$ is a factor of the word $[t_\ell]$, whence $[t_\ell]$ contains a factor of the form (7.2b) or (7.2d), depending on whether or not the letters x_1, x_2 , and q are distinct. But this is impossible by (ii) of Lemma 7.5. Therefore,

(B) the word $[y_i \varphi]$ is a singleton.

Suppose that $c_i = 1$ and $[h_i^{c_i} \varphi]$ is non-singleton. Then, $\overline{[h_i^{c_i} \varphi]} = \dots x_1 x_2 \dots$ for some $x_1, x_2 \in \mathcal{A}$. Choose any $q \in \text{con}(\overline{[x\varphi]})$ and $t \in \text{con}(\overline{[y_i \varphi]})$. Then,

$$\begin{aligned} [W^\uparrow \varphi] &= [x\varphi] \dots [y_i \varphi][h_i^{c_i} \varphi][(y_i \varphi)^*] \dots [(x\varphi)^*] \\ &= \underbrace{(\dots q^{\otimes_1} \dots)}_{[x\varphi]} \dots \underbrace{(\dots t^{\otimes_2} \dots)}_{[y_i \varphi]} \underbrace{(\dots x_1^{\otimes_3} x_2^{\otimes_4} \dots)}_{[h_i^{c_i} \varphi]} \underbrace{(\dots t^{\otimes_5} \dots)}_{[(y_i \varphi)^*]} \dots \underbrace{(\dots q^{\otimes_6} \dots)}_{[(x\varphi)^*]} \end{aligned}$$

for some $\otimes_j \in \{1, *\}$. Since $W^\uparrow \varphi$ is a subterm of t_ℓ , by (iii) of Remark 7.1, either $[W^\uparrow \varphi]$ or $[(W^\uparrow \varphi)^*]$ is a factor of the word $[t_\ell]$, whence $[t_\ell]$ contains a subterm of the form (7.2b), (7.2d), or (7.2e), depending on whether or not the letters x_1, x_2, q , and t are distinct. But this is impossible by (ii) of Lemma 7.5. Therefore,

(C) if $c_i = 1$, then the word $[h_i^{c_i} \varphi]$ is a singleton.

Now since $\mathbf{u}_\ell \varphi = \bar{w}^\uparrow \varphi$ is a subterm of \mathbf{t}_ℓ , by (iii) of Remark 7.1, either $\llbracket w^\uparrow \varphi \rrbracket$ or $\llbracket (w^\uparrow \varphi)^* \rrbracket$ is a factor of the word $\llbracket \mathbf{t}_\ell \rrbracket = v_{\text{fix}}^\uparrow$, where

$$\llbracket w^\uparrow \varphi \rrbracket = \llbracket x\varphi \rrbracket \left(\prod_{i=1}^n (\llbracket y_i \varphi \rrbracket \llbracket h_i^c \varphi \rrbracket \llbracket (y_i \varphi)^* \rrbracket) \right) \llbracket (x\varphi)^* \rrbracket$$

and

$$\llbracket (w^\uparrow \varphi)^* \rrbracket = \llbracket x\varphi \rrbracket \left(\prod_{i=n}^1 (\llbracket y_i \varphi \rrbracket \llbracket (h_i^c \varphi)^* \rrbracket \llbracket (y_i \varphi)^* \rrbracket) \right) \llbracket (x\varphi)^* \rrbracket.$$

Therefore, (A)–(C) imply that $m = n$ and either $v_{\text{fix}}^\uparrow = \llbracket w^\uparrow \varphi \rrbracket$ or $v_{\text{fix}}^\uparrow = \llbracket (w^\uparrow \varphi)^* \rrbracket$. It follows that $v_{\text{fix}}^\uparrow \approx v_{\text{fix}}^\downarrow$ coincides with the identity $w^\uparrow \approx w^\downarrow$ and so also the identity $\mathbf{u}_\ell \approx \mathbf{v}_\ell$. \square

Acknowledgement: The author would like to thank the anonymous referees for helpful suggestions, and Igor Dolinka and Marcel G. Jackson for very fruitful discussions.

References

- [1] Y. A. Bahturin and A. Y. Ol'shanskiĭ, Identical relations in finite Lie rings, *Math. USSR Sb.* **25** (1975), 507–523.
- [2] G. Birkhoff, On the structure of abstract algebra, *Proc. Cambridge Philos. Soc.* **31** (1935), 433–454.
- [3] J. Hage and T. Harju, On involutions arising from graphs, in: *Algorithmic Bioprocesses*, Nat. Comput. Ser., Springer, Berlin (2009), 623–630.
- [4] M. Jackson, Finite semigroups with infinite irredundant identity bases, *Internat. J. Algebra Comput.* **15** (2005), 405–422.
- [5] J. Ježek, Nonfinitely based three-element idempotent groupoids, *Algebra Universalis* **20** (1985), 292–301.
- [6] J. Kad'ourek, On varieties of combinatorial inverse semigroups. II, *Semigroup Forum* **44** (1992), 53–78.
- [7] R. L. Kruse, Identities satisfied by a finite ring, *J. Algebra* **26** (1973), 298–318.
- [8] E. W. H. Lee, A sufficient condition for the non-finite basis property of semigroups, *Monatsh. Math.* **168** (2012), 461–472.
- [9] E. W. H. Lee, A class of finite semigroups without irredundant bases of identities, to appear in *Yokohama Math. J.*
- [10] E. W. H. Lee, Finitely based finite involution semigroups with non-finitely based reducts, to appear in *Quaest. Math.*
- [11] E. W. H. Lee, J. R. Li and W. T. Zhang, Minimal non-finitely based semigroups, *Semigroup Forum* **85** (2012), 577–580.
- [12] I. V. L'vov, Varieties of associative rings. I, *Algebra Logic* **12** (1973), 150–167.
- [13] G. I. Mashevitskiĭ, An example of a finite semigroup without an irreducible basis of identities in the class of completely 0-simple semigroups, *Russian Math. Surveys* **38** (1983), no. 2, 192–193.
- [14] R. McKenzie, Tarski's finite basis problem is undecidable, *Internat. J. Algebra Comput.* **6** (1996), 49–104.
- [15] V. L. Murskiĭ, The existence in the three-valued logic of a closed class with a finite basis having no finite complete system of identities, *Soviet Math. Dokl.* **6** (1965), 1020–1024.
- [16] S. Oates and M. B. Powell, Identical relations in finite groups, *J. Algebra* **1** (1964), 11–39.
- [17] P. Perkins, Bases for equational theories of semigroups, *J. Algebra* **11** (1969), 298–314.
- [18] M. V. Sapir, On Cross semigroup varieties and related questions, *Semigroup Forum* **42** (1991), 345–364.
- [19] M. V. Volkov, The finite basis problem for finite semigroups, *Sci. Math. Jpn.* **53** (2001), 171–199.

Received May 29, 2014; revised January 3, 2015.