

8-2016

An Update on a Few Permanent Conjectures

Fuzhen Zhang

Nova Southeastern University, zhang@nova.edu

Follow this and additional works at: https://nsuworks.nova.edu/math_facarticles

 Part of the [Discrete Mathematics and Combinatorics Commons](#)

NSUWorks Citation

Zhang, Fuzhen, "An Update on a Few Permanent Conjectures" (2016). *Mathematics Faculty Articles*. 211.
https://nsuworks.nova.edu/math_facarticles/211

This Article is brought to you for free and open access by the Department of Mathematics at NSUWorks. It has been accepted for inclusion in Mathematics Faculty Articles by an authorized administrator of NSUWorks. For more information, please contact nsuworks@nova.edu.

Research Article

Open Access

Special Issue: 5th International Conference on Matrix Analysis and Applications

Fuzhen Zhang

An update on a few permanent conjectures

DOI 10.1515/spma-2016-0030

Received March 18, 2016; accepted July 27, 2016

Abstract: We review and update on a few conjectures concerning matrix permanent that are easily stated, understood, and accessible to general math audience. They are: Soules permanent-on-top conjecture[†], Lieb permanent dominance conjecture, Bapat and Sunder conjecture[†] on Hadamard product and diagonal entries, Chollet conjecture on Hadamard product, Marcus conjecture on permanent of permanents, and several other conjectures. Some of these conjectures are recently settled; some are still open. We also raise a few new questions for future study. ([†]conjectures have been recently settled negatively.)

Keywords: Bapat-Sunder conjecture; Chollet conjecture; Drury conjecture; Foregger conjecture; Liang-So-Zhang conjecture; Lie conjecture; Marcus conjecture; Marcus-Minc conjecture; permanent; permanent dominance conjecture; permanent-on-top conjecture; Schur power matrix

MSC: 15A15

In memory of Professors Marvin Marcus (1927-2016) and Ingram Olkin (1924-2016).

1 Introduction

Computed from the elements of a square matrix, the determinant of a square matrix is one of the most useful and important concepts in mathematics. Of many matrix functions, *permanent* is another important one. It arises naturally in the study of the symmetric tensors in multilinear algebra (see, e.g., [37]); it also plays a role in combinatorics (see, e.g., [56]). Both terms were introduced in the 1800s (see, e.g., [39, p. 1]); and they are still useful in research.

One of the intriguing problems on permanent is the so-called van der Waerden conjecture (theorem) regarding the minimum value of the permanent on the Birkhoff polytope of doubly stochastic matrices. It was conjectured by van der Waerden in 1926 and resolved in the affirmative by Egoryčev and Falikman independently in 1981 (see, e.g., [40]). Second to the van der Waerden conjecture is, in my opinion, the *permanence dominance conjecture* (see below). Since its appearance in the mid-1960s, it has drawn much attention of the mathematicians in the area; it remains open as one of the most important unsolved problems in linear algebra and matrix theory. A closely related and stronger statement is the *permanent-on-top* conjecture which appeared about the same time and is recently shown to be false with the help of computation utility.

We begin by the permanent-on-top (POT) conjecture and two chains of conjectures on permanent starting with the POT conjecture, reviewing and updating their developments and status. We also include a few other conjectures on permanent of our interest that are easily stated but remain unsolved or even have no progress over the decades. A few comprehensive surveys on permanent, including many open conjectures and problems, are the Minc's monograph *Permanents* 1978 [39], and subsequent articles *Theory of Permanents 1978–1981* [40] and *Theory of Permanents 1982–1985* [41], followed by Cheon and Wanless' *An update on Minc's survey of open problems involving permanents* 2005 [9].

Fuzhen Zhang: Department of Mathematics, Halmos College, Nova Southeastern University, 3301 College Ave, Fort Lauderdale, FL 33314, USA, E-mail: zhang@nova.edu

2 Preliminaries

Let M_n be the set of all $n \times n$ complex matrices. We assume $n \geq 2$ in the paper. For an $n \times n$ matrix $A = (a_{ij})$, the determinant $\det A$ of A is $\sum_{\sigma \in S_n} \prod_{t=1}^n \text{sign}(\sigma) a_{t\sigma(t)}$, where S_n is the symmetric group of degree n , while the permanent of $A = (a_{ij})$, denoted by $\text{per } A$, is simply defined as $\text{per } A = \sum_{\sigma \in S_n} \prod_{t=1}^n a_{t\sigma(t)}$. For $A \in M_n$, by writing $A > 0$ (resp. $A \geq 0$) we mean that A is positive (resp. semi) definite, that is, $x^*Ax > 0$ (resp. $x^*Ax \geq 0$) for all column vectors $x \in \mathbb{C}^n$, where $x^* = (\bar{x})^t$ is the conjugate transpose of x . It is known that if A is positive semidefinite, then $0 \leq \det A \leq \prod_{i=1}^n a_{ii} \leq \text{per } A$. Determinant and permanent may be regarded as sister functions of positive semidefinite matrices as many of their results exist side by side. For example, when A and B are $n \times n$ and positive semidefinite, then $\det(A+B) \geq \det A + \det B$ and $\text{per}(A+B) \geq \text{per } A + \text{per } B$. (Note: the directions of inequalities for “det” and “per” sometimes are the same and some other times are reversed.) A permanent analogue of a determinant result does not always exist. For instance, if $A \geq 0$ is written as $A = R + Si$, where R and S are Hermitian, then $\det A \leq \det R$ (the Robertson-Taussky inequality); but $\text{per } A$ and $\text{per } R$ are incomparable in general [27].

Determinant and permanent are special *generalized matrix functions*. Let S_n be the symmetric group of degree n , H be a subgroup of S_n , and χ be a character on H . The generalized matrix function of an $n \times n$ matrix $A = (a_{ij})$ with respect to H and χ is defined by (see, e.g., [30, p. 124])

$$d_\chi^H(A) = \sum_{\sigma \in H} \chi(\sigma) \prod_{t=1}^n a_{t\sigma(t)}$$

Setting $H = S_n$ and $\chi(g) = \text{sign}(g) = \pm 1$ according to $g \in H$ being even or odd, we have the determinant $\det A$; putting $H = S_n$ and $\chi(g) = 1$ for all g , we get the permanent $\text{per } A = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i\sigma(i)}$. The product of the main diagonal entries of A , $h(A) = a_{11}a_{22} \cdots a_{nn}$, known as the Hadamard matrix function, is also a generalized matrix function by taking $H = \{e\}$, where e is the group identity of S_n . Specially, if $H = S_n$ and χ is an irreducible character of S_n , then the permanent $d_\chi^H(A)$ is referred to as *immanent*.

Let $A = (a_{ij})$ be a square matrix. We write $A = (A_{ij}) \in M_n(M_m)$ if A is an $n \times n$ partitioned matrix in which each block A_{ij} is $m \times m$. (Thus, A is an $mn \times mn$ matrix.) In case $n = 2$, we write explicitly $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}$.

Listed below are a few well-known milestone inequalities for determinant, permanent, and generalized matrix functions of positive semidefinite matrices. They stand out because of their elegance in form and importance in applications. Let $A = (a_{ij})$ be an $n \times n$ positive semidefinite matrix with a partitioned form $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}$, where A_{11} and A_{22} are square submatrices (possibly of different sizes) of A . We have the following inequalities concerning A .

Hadamard inequality (1893):

$$\det A \leq \prod_{i=1}^n a_{ii} \quad (1)$$

Fischer inequality (1908):

$$\det A \leq \det A_{11} \det A_{22} \quad (2)$$

Schur inequality (1918):

$$\det A \leq \frac{1}{\chi(e)} d_\chi^H(A) \quad (3)$$

Marcus inequality (1963):

$$\text{per } A \geq \prod_{i=1}^n a_{ii} \quad (4)$$

Lieb inequality (1966):

$$\text{per } A \geq \text{per } A_{11} \text{ per } A_{22} \quad (5)$$

Schur inequality (3) implies Fischer inequality (2) which implies Hadamard inequality (1). (4) and (5) are permanent analogues of (1) and (2), respectively. Inequality (5) (in a more general form) was first conjectured by Marcus and Newman in 1965 [32] and proved by Lieb in 1966 [28], and re-proved by Djoković in 1969 [11].

If we compare (1) and (4), and (2) and (5), in view of (3), we can naturally ask if the right hand of (3) is bounded by $\text{per } A$, that is, if the following holds:


$$\frac{1}{\chi(e)} a_\chi^H(A) \leq \text{per } A \tag{6}$$

This is a conjecture of Lieb also known as the *permanent dominance conjecture*. It was stated explicitly by Lieb in 1966 [28] when studying a similar problem of Marcus and Minc 1965 [32]: Under what conditions on χ and H will the following inequality hold for all $n \times n$ positive semidefinite A

$$\sum_{\sigma \in H} \chi(\sigma) \prod_{i=1}^n a_{i\sigma(i)} \leq \text{per } A? \tag{7}$$

(6) and (7) are the same when χ is degree 1, i.e., $\chi(e) = 1$.

A great amount of effort was made in attempting to solve the Lieb permanent dominance conjecture, especially from the late 1960s to early 1990s. Motivated by Lieb conjecture [28], Soules proposed in his 1966 Ph.D. thesis (see also Merris 1987 [36]) a conjecture stronger than the permanent dominance conjecture. Soules conjecture states that the permanent of a positive semidefinite matrix A is the largest eigenvalue of the Schur power matrix of A (see Section 3). This is referred to as permanent-on-top (POT) conjecture. In fact, POT conjecture is the strongest among several permanent conjectures (see, e.g., [9]):



It is known now that Soules and Bapat & Sunder conjectures are false; but the other three, the Lieb, Marcus and Chollet conjectures, are still open. In view of recent developments in the area and with more advanced computation tools available nowadays, it is worth revisiting these long standing conjectures.

3 The permanent-on-top conjecture is false

In a recent publication, Shchesnovich 2016 [52] settled in the negative a long standing conjecture on permanent; it shows that the permanent-on-top (POT) conjecture is false. The POT conjecture was originally formulated by Soules in his Ph. D. dissertation 1966 [53, Conjecture, p. 3] (formally published in Minc’s 1983 [40, p. 249]) in attempting to answer the Lieb conjecture [28]. It stated that the permanent of a positive semidefinite matrix A was the largest eigenvalue of the Schur power matrix of A . For an $n \times n$ matrix $A = (a_{ij})$, the *Schur power matrix* of A , denoted by $\pi(A)$, is an $n! \times n!$ matrix with entries $s_{\alpha\beta}$ lexicographically indexed by permutations $\alpha, \beta \in S_n$:

$$s_{\alpha\beta} = \prod_{t=1}^n a_{\alpha(t)\beta(t)}$$

As a principal submatrix of the n -fold tensor (Kronecker) power $\otimes^n A$, the Schur power matrix $\pi(A)$ inherits many properties of A . For instance, if A is positive semidefinite, then so is its Schur power; and the eigenvalues of $\pi(A)$ interlace those of $\otimes^n A$. Observing that for each fixed $\alpha \in S_n$,

$$\sum_{\beta \in S_n} s_{\alpha\beta} = \sum_{\beta \in S_n} \prod_{t=1}^n a_{\alpha(t)\beta(t)} = \text{per } A$$

We see that every row sum of $\pi(A)$ is $\text{per } A$ [54]. Consequently, $\text{per } A$ is an eigenvalue of $\pi(A)$. It is also known (and not difficult to show) that $\det A$ is an eigenvalue of $\pi(A)$ too. In fact, $\det A$ is the smallest eigenvalue of $\pi(A)$ when $A \geq 0$; see [37, p. 221]. For positive semidefinite A , the POT conjecture claimed that $\text{per } A$ was the largest eigenvalue of $\pi(A)$. Note that POT conjecture is independent of subgroup H and character χ in (6).

The counterpart of the POT conjecture for determinant is that $\det A$ is the smallest eigenvalue of $S(A)$ which is shown implicitly in Schur 1918 [51], as pointed out by Bapat and Sunder 1986 [4, p. 154]. Bapat and Sunder 1986 [4] proved that the POT conjecture is true for $n \leq 3$ (and also provided an equivalent form of the POT conjecture). POT conjecture involved no subgroup and character and it would yield Lieb conjecture (in the case of degree 1 characters as assumed in [53, p. 3] and more generally in Merris 1987 [36]) because the value on the left hand side of (6) is contained in the numerical range of the Schur power matrix. If $W(X)$ denotes the numerical range of a square matrix X , then the POT conjecture is equivalent to the statement that for $A \geq 0$,

$$W(\pi(A)) = [\det A, \text{per } A] \quad (8)$$

In fact, Soules POT conjecture was the strongest among several permanent conjectures; see the conjecture chains in the previous section; see also, e.g., [9].

Notice that if A is an $n \times n$ positive semidefinite matrix, then $x^*Ax \leq \lambda_{\max}(A)x^*x$ for all column vectors $x \in \mathbb{C}^n$, where $\lambda_{\max}(A)$ is the largest eigenvalue of A . Shchesnovich 2016 [52] presented an example of 5×5 positive semidefinite matrix H (thus $\pi(H)$ is $5! \times 5!$, i.e., 120×120) and a column vector X of $5! = 120$ components (which is omitted here as it is too large) such that

$$X^* \pi(H)X > \text{per}(H)X^*X$$

where

$$H = \begin{pmatrix} 40 & 14 - 22i & -4 - 8i & 8 + 16i & 22 - 36i \\ 14 + 22i & 22 & 16 - 14i & -9 - i & 23 - 12i \\ -4 + 8i & 16 + 14i & 52 & 8 - 34i & 14 - 30i \\ 8 - 16i & -9 + i & 8 + 34i & 34 & 18 - 19i \\ 22 + 36i & 23 + 12i & 14 + 30i & 18 + 19i & 75 \end{pmatrix} = u^*u + v^*v$$

with

$$u = (4 + 2i, 2 - 3i, -4 - 4i, -3 + 4i, 1), \quad v = (2 - 4i, -3i, 2 - 4i, -3i, -5 - 7i).$$

H is a 5×5 positive semidefinite matrix of rank 2 having eigenvalues 0, 0, 0, 91, and 132, while its Schur power matrix $\pi(H)$ is of rank 27.

The Schur power matrix $S(H)$ is of order $5! = 120$. It would be difficult to compute and display $\pi(H)$. A computation utility is needed. It is also demonstrated in [52] through characteristic polynomial that $\lambda_{\max}(\pi(H)) > \text{per } H$:

$$\lambda_{\max}(\pi(H)) = 320(2185775 + \sqrt{160600333345}) > \text{per}(H) = 814016640$$

For an n -square matrix A , the Schur power matrix $\pi(A)$ as a principal submatrix of the tensor power $\otimes^n A$ may have interest in its own right. It is easy to show that $\{w_1 w_2 \cdots w_n \mid w_i \in W(A), i = 1, 2, \dots, n\} \subseteq W(\otimes^n A)$.

Question 1: Can one find a counterexample of size $n = 4$ for the POT conjecture? As is known, the POT conjecture is true for $n \leq 3$. With a counterexample of 5×5 , Shchesnovich 2016 [52, p. 198] states that a 4×4 counterexample was not found despite an extensive search.

Question 2: Referring (8), what is exactly the maximum value (which may exceed $\text{per } A$) in the numerical range of the Schur power matrix of a positive semidefinite matrix? Of a Hermitian matrix? That is, find $\lambda_{\max}(\pi(A))$. This question is especially of interest as the numerical range of a Hermitian matrix is a closed interval $[d, p]$. What are the left and right end points, i.e. d and p , of the interval? A result of Pate 1989 [46, Theorem 4] implies that $\langle \pi(A)x, x \rangle \leq \text{per } A$ for all unit vectors x in certain (but not all) subspaces of $\mathbb{C}^{n!}$.

Question 3: Investigate the numerical range of the Schur power matrix $\pi(A)$ for a general n -square matrix A . It is contained in the numerical range of $\otimes^n A$ because $\pi(A)$ is a principal submatrix of the latter.

Note: There has been a confusion about the names of the conjectures in the literature. In 1987 [36, Conjecture 1], Merris called inequality (6) the *Permanental Dominance Conjecture*; in the same book, Johnson 1987 [26, p. 168] called (6) the *permanent-on-top conjecture*, while in their introductions, Bapat [2] and Zhang [58] considered the two being the same. Besides, there have been several variations of the POT conjecture; see Merris 1987 [36].

Now it is known that the Soules POT conjecture is false. The weaker one, Lieb permanence dominance conjecture, is still open. It has been proven true for special cases of $H = S_n$ and some characters χ (see the next section).

4 The permanent dominance conjecture is open

In 1965, Marcus and Minc proposed the following question regarding the permanent of positive semidefinite matrices with a subgroup of the permutation group and group character (see also [39, p. 158], [40, p. 244], and [41, p. 132]):

Problem 2 of [32, p. 588]. *Let H be a subgroup of S_n and let χ be a character of degree 1 of H . Under what conditions on χ and H will the inequality*

$$\sum_{\sigma \in H} \chi(\sigma) \prod_{i=1}^n a_{i\sigma(i)} \leq \text{per } A \quad (9)$$

hold for all $n \times n$ positive semidefinite Hermitian A ?

In 1966, Lieb stated that the answer to the above question is given by the conjecture below (referred to as Conjecture α in Lieb 1966 [28]):

Lieb permanent dominance conjecture 1966 [28]. *Let H be a subgroup of the symmetric group S_n and let χ be a character of degree m of H . Then*

$$\frac{1}{m} \sum_{\sigma \in H} \chi(\sigma) \prod_{i=1}^n a_{i\sigma(i)} \leq \text{per } A \quad (10)$$

holds for all $n \times n$ positive semidefinite Hermitian A .

The permanent dominance conjecture (10) claims that $\text{per } A$ is the smallest (best) upper-bound of $\frac{1}{m} \sum_{\sigma \in H} \chi(\sigma) \prod_{i=1}^n a_{i\sigma(i)}$ for all subgroups and characters. Merris 1983 [34] gave an upper bound

$$\frac{1}{m} \sum_{\sigma \in H} \chi(\sigma) \prod_{i=1}^n a_{i\sigma(i)} \leq \left(\prod_{i=1}^n (A^n)_{ii} \right)^{1/n} \leq \frac{1}{n} \sum_{i=1}^n \lambda_i^n \quad (11)$$

where $(A^n)_{ii}$ denotes the i th diagonal entry of A^n and the λ_i s are the eigenvalues of A . The middle term in (11) is no less than $\text{per } A$ as noted in [36, p. 215].

Historically, in his Ph.D. dissertation, Soules 1966 [54, p. 3] (where characters were assumed to have degree 1) proposed the later so-called POT conjecture, stating that it had the permanent dominance inequality as its consequence. This method for approaching (9) and (10) was announced by Soules at the 1981 Santa Barbara Multilinear Algebra Conference (see [40, p. 245]).

In 1986 at the Auburn Matrix Conference, Merris called Lieb's conjecture *Permanent Dominance Conjecture* (see [35, p. 135], and also [36]). This term has later been used by Minc [41, p. 135], Soules [54], Pate [49], Cheon and Wanless [9], and many others. So it has become known as the *permanent dominance conjecture*. Merris [36, p. 216] explicitly pointed out that the Soules POT conjecture implied the Lieb permanent dominance conjecture. In his 1994 paper [54], Soules gave a clear and brief explanation of this. In fact, Soules POT

conjecture would imply several permanent conjectures (see, e.g., [9]). A great amount of work was devoted to the permanent dominance conjecture since its appearance to early 1990s. As an open problem on permanent, this conjecture is arguably most interesting and intriguing; it involves group theory, combinatorics, and matrix theory, of course.

Several paths have been taken to attack or approach the Lieb conjecture. Observe that the right hand side of (10) is independent of the choices of the subgroup and the character. Immediate cases to be studied are: $H = S_n$, χ is the principal character (i.e., $\chi(e) = 1$), or χ is irreducible.

Great amount work has been done for immanant (with $H = S_n$ and irreducible χ) and some significant results have been obtained. It is known [48, 49] that the permanent dominance conjecture is true for immanants with $n \leq 13$.

The interest and attention on this interesting and rather difficult conjecture appeared fading away in the last decades. In the author’s opinion, there is still a long way to go for a complete solution of the general conjecture. Cheon and Wanless 2005 [9, p. 317] set an excellent account in detail about the permanent dominance conjecture. Other references on this conjecture are Merris 1987 [36], Johnson 1987 [26], and references therein.

5 Bapat & Sunder conjecture is false

Shchesnovich 2016 [52] devotes a section discussing a permanent conjecture of Bapat and Sunder 1985 [3] that is weaker than the POT conjecture and explains that the weaker one may be true for a physical reason (quantum theory). The counterexample provided in [52] settles POT conjecture negatively but does not work for the Bapat & Sunder conjecture.

Recall the Oppenheim inequality [42] for positive semidefinite matrices: Let $A = (a_{ij})$ and $B = (b_{ij})$ be $n \times n$ positive semidefinite matrices. Then

$$\det(A \circ B) \geq \det A \prod_{i=1}^n b_{ii} \geq \det A \det B \tag{12}$$

where $A \circ B = (a_{ij}b_{ij})$ is the Hadamard (a.k.a Schur) product of A and B . In view of (12) for permanent, Bapat and Sunder 1985 [3, p. 117] and 1986 [4, Conjecture 2] raised the question (conjecture) whether

$$\text{per}(A \circ B) \leq \text{per } A \, b_{11} \cdots b_{nn} \tag{13}$$

It is not difficult to reduce Bapat & Sunder conjecture to the case of correlation matrices (see, e.g., [57]). By a *correlation matrix*, we mean a positive semidefinite matrix all whose main diagonal entries are equal to 1. That is, Bapat & Sunder conjecture is equivalent to the statement that if A and B are correlation matrices, then

$$\text{per}(A \circ B) \leq \text{per } A \tag{14}$$

Bapat & Sunder conjecture is weaker than the permanent-on-top conjecture (which is now known false); see Merris 1987 [36, Conjecture 10, p. 220]. Drury 2016 [14] found a counterexample and settled this negatively. Let A be

$$\begin{pmatrix} 1 & 0 & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ 0 & 1 & \frac{1}{\sqrt{2}}e^{\frac{4}{5}in} & \frac{1}{\sqrt{2}}e^{\frac{2}{5}in} & \frac{1}{\sqrt{2}}e^{-\frac{2}{5}in} & \frac{1}{\sqrt{2}}e^{-\frac{4}{5}in} & \frac{1}{\sqrt{2}}e^{-\frac{4}{5}in} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}}e^{-\frac{4}{5}in} & 1 & \cos\left(\frac{1}{5}\pi\right)e^{-\frac{1}{5}in} & \cos\left(\frac{2}{5}\pi\right)e^{-\frac{2}{5}in} & \cos\left(\frac{2}{5}\pi\right)e^{\frac{2}{5}in} & \cos\left(\frac{1}{5}\pi\right)e^{\frac{1}{5}in} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}}e^{-\frac{2}{5}in} & \cos\left(\frac{1}{5}\pi\right)e^{\frac{1}{5}in} & 1 & \cos\left(\frac{1}{5}\pi\right)e^{-\frac{1}{5}in} & \cos\left(\frac{2}{5}\pi\right)e^{-\frac{2}{5}in} & \cos\left(\frac{2}{5}\pi\right)e^{\frac{2}{5}in} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}}e^{\frac{2}{5}in} & \cos\left(\frac{2}{5}\pi\right)e^{\frac{2}{5}in} & \cos\left(\frac{1}{5}\pi\right)e^{\frac{1}{5}in} & 1 & \cos\left(\frac{1}{5}\pi\right)e^{-\frac{1}{5}in} & \cos\left(\frac{2}{5}\pi\right)e^{-\frac{2}{5}in} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}}e^{\frac{2}{5}in} & \cos\left(\frac{2}{5}\pi\right)e^{-\frac{2}{5}in} & \cos\left(\frac{2}{5}\pi\right)e^{\frac{2}{5}in} & \cos\left(\frac{1}{5}\pi\right)e^{\frac{1}{5}in} & 1 & \cos\left(\frac{1}{5}\pi\right)e^{-\frac{1}{5}in} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}}e^{\frac{4}{5}in} & \cos\left(\frac{1}{5}\pi\right)e^{-\frac{1}{5}in} & \cos\left(\frac{2}{5}\pi\right)e^{-\frac{2}{5}in} & \cos\left(\frac{2}{5}\pi\right)e^{\frac{2}{5}in} & \cos\left(\frac{1}{5}\pi\right)e^{\frac{1}{5}in} & 1 \end{pmatrix}$$

Drury’s matrices A (of rank 2 with complex entries) and $B = A^t = \bar{A}$ are 7×7 correlation matrices with $\text{per } A = 45$ and $\text{per}(A \circ B) = \frac{6185}{128}$, having the ratio $\text{per}(A \circ B) / (\text{per } A \prod_{i=1}^7 b_{ii}) = 1237/1152 > 1$. Moreover, $\lambda_{\max}(A) = 3.5$, while $\lambda_{\max}(\pi(A)) = 525/8 = 65.625$. Very recently, Drury [15] provided a *real* counterexample of size 16×16 with rank 3. So $n = 16$ is the smallest size of known real positive semidefinite matrices that fail the POT conjecture. Note that Drury’s counterexamples also disprove the POT conjecture as the latter is stronger.

Question 4: Can one find a counterexample of size $n < 7$, say $n = 4$, with complex (or real) entries, for the Bapat & Sunder conjecture?

Question 5: Can one find a *real* counterexample of size $n < 16$? That is to say, does the Bapat & Sunder conjecture (and also the POT conjecture) hold true for *real* positive semidefinite matrices of sizes no more than 15? Shchesnovich 2016 [52, p. 198] states that all found counterexamples are complex and rank deficient. Drury’s counterexamples are also rank deficient. Soules 1994 [54] showed certain necessary conditions for a *real* positive semidefinite matrix to fail the conjecture. (Note: the statement “By Theorem 1, the conjecture can only fail at a singular matrix” on page 222, line 19 of [54] is incorrect.) In particular, for the case of real and $n = 4$, if POT fails, then there exists a *singular* matrix for which the POT is false. It follows that the POT conjecture would be true for $n = 4$ if one could prove that it is true for all *singular* positive semidefinite matrices of size $n = 4$.

Denote by \mathcal{C}_n the collection of all $n \times n$ complex correlation matrices. The set \mathcal{C}_n can be thought of as a subset of \mathbb{C}^{n^2} and it is compact and convex. The compactness of \mathcal{C}_n follows from the fact that $A \geq 0$ if and only if all principal submatrices of A have nonnegative determinants. We also see that \mathcal{C}_n is closed under the Hadamard product. Given $A \in \mathcal{C}_n$, we define a function on \mathcal{C}_n by

$$f_A(X) = \text{per}(A \circ X), \quad X \in \mathcal{C}_n \tag{15}$$

Since \mathcal{C}_n is compact and f_A is continuous, there exists a correlation matrix depending on A , referred to as *maximizer* of A and denoted by M_A , such that

$$f(M_A) = \max_{X \in \mathcal{C}_n} \text{per}(A \circ X)$$

i.e., the maximal value can be attained. Drury’s example shows that it is possible that a (irreducible) maximizer has all entries with moduli less than 1. Several properties of maximizing matrices are presented in Zhang 2013 [58].

In view of (12), we see that the determinant is a “Hadamard-dilation” function on \mathcal{C}_n in the sense that $\det A \leq \det(A \circ X)$ for $A \in \mathcal{C}_n$ and any $X \in \mathcal{C}_n$. Equivalently, $\max\{\det A, \det B\} \leq \det(A \circ B)$ for $A, B \in \mathcal{C}_n$. In contrast, the permanent is no “Hadamard-compression” on \mathcal{C}_n .




Figure 1: \det , per , and a_{ij}^H on \mathcal{C}_n

Let $A = (a_{ij}) \in M_n$ and k be a positive integer. Denote by $A^{[k]}$ the k -fold Hadamard product of A , that is, $A^{[k]} = (a_{ij}^k)$. If A is a correlation matrix, then $\lim_{k \rightarrow \infty} A^{[k]}$ is a correlation matrix having entries 1, -1 , and 0.

It is easy to see that $\text{per}(A^{[k]}) \leq \text{per} A$ if k is large enough. If $A \in \mathbb{C}_n$ is such a matrix that has an maximizer M_A for which $\text{per}(A \circ M_A) > \text{per} A$, then there must exist a positive integer r such that $\text{per}(A \circ M_A^{[r]}) \leq \text{per} A$. Drury’s 16×16 example shows that $\text{per}(A \circ A) > \text{per} A$ for some real correlation matrices A .

Question 6: Characterize all the matrices $A \in \mathbb{C}_n$ that have the Hadamard-compression property, that is, $\text{per}(A \circ X) \leq \text{per} A$ for all $X \in \mathbb{C}_n$. These are exactly the matrices having the property $\text{per}(A \circ M_A) \leq \text{per} A$.

Question 7: Other than the determinant, are there any other generalized matrix functions d_X^H on \mathbb{C}_n that are Hadamard-dilation? Namely, for all $n \times n$ correlation matrices A and B , $d_X^H(A \circ B) \geq \max\{d_X^H(A), d_X^H(B)\}$. Note that such a matrix function is closely related to the subgroup H and character. If $H = \{e\}$ (the Hadamard matrix function), then the inequality holds true.

Question 8: Characterize all maximizers of a given matrix $A \in \mathbb{C}_n$.

Moreover, some other fragmentary results are seen. It is known [57, Result 5] that for $n \times n$ correlation matrices A and B , $\text{per}(A \circ B) \leq \text{per} C_t$, where C_t is the $n \times n$ correlation matrix with all off-diagonal entries equal to t , where $t = 1 - \lambda_{\min}(A)$, $t = 1 - \lambda_{\min}(B)$, or $t = 1 - \lambda_{\min}(A \circ B)$. It is also known [57, Result 4] that if A is nonnegative entrywise, i.e., $a_{ij} \geq 0$, and B is positive semidefinite, then $|\text{per}(A \circ B)| \leq \text{per}(A)b_{11} \cdots b_{nn}$.

6 Chollet conjecture is open

In view of the classical result (of Oppenheim 1930) $\det(A \circ B) \geq \det A \det B$ for $A, B \geq 0$, Chollet 1982 [10] proposed an interesting problem concerning the permanent of the Hadamard product of positive semidefinite matrices:

Chollet conjecture 1982 [10]. If $A \geq 0$ and $B \geq 0$ are $n \times n$ matrices, then

$$\text{per}(A \circ B) \leq \text{per} A \text{ per} B \tag{16}$$

Chollet showed that the inequality holds if and only if it is true when $B = \bar{A}$:

$$\text{per}(A \circ \bar{A}) \leq (\text{per} A)^2 \tag{17}$$

(17) is immediate from (16) with $B = \bar{A}$. (17) implies (16) as shown in [10]:

$$\begin{aligned} 0 \leq \text{per}(A \circ B) &= \sum_{\sigma \in S_n} \prod_{t=1}^n a_{t\sigma(t)} b_{t\sigma(t)} \\ &\leq \sum_{\sigma \in S_n} \prod_{t=1}^n |a_{t\sigma(t)}| |b_{t\sigma(t)}| \\ &\leq \left(\sum_{\sigma \in S_n} \prod_{t=1}^n |a_{t\sigma(t)}|^2 \right)^{\frac{1}{2}} \left(\sum_{\sigma \in S_n} \prod_{t=1}^n |b_{t\sigma(t)}|^2 \right)^{\frac{1}{2}} \\ &= (\text{per}(A \circ \bar{A}))^{\frac{1}{2}} (\text{per}(B \circ \bar{B}))^{\frac{1}{2}} \\ &\leq \text{per} A \text{ per} B \end{aligned} \tag{18}$$

With Drury’s example in [14], $\text{per}(A \circ B) = \text{per}(A \circ \bar{A}) = \frac{6185}{128} < 50 < 45^2 = (\text{per} A)^2$. Moreover, if $A = (a_{ij})$ is a correlation matrix, then all $|a_{ij}| \leq 1$. Thus, $\text{per}(A \circ \bar{A}) \leq \text{per}(|A|)$, where $|A| = (|a_{ij}|)$. Note that $\text{per} A \leq \text{per}(|A|)$.

Chollet conjecture is weaker than the Bapat & Sunder conjecture because $\text{per} B \geq \prod_{i=1}^n b_{ii}$. Special cases of (16) were settled in the 1980s. Gregorac and Hentzel 1987 [17] showed by elementary methods that the inequality is true for the case of 2×2 and 3×3 matrices, which are also immediate consequences of the stronger results of Bapat and Sunder 1986 [4]. Grone and Merris 1987 [20] and also Marcus and Sandy 1988

[33] discussed the Chollet conjecture and the Bapat & Sunder conjecture. Related works are seen in Zhang 1989 [57] and Beasley 2000 [5] (on some infinite classes of matrices). As the Bapat & Sunder conjecture (13) is settled in the negative recently, Chollet conjecture stands out and is appealing as an interesting open question. The real case of the Chollet conjecture is the following.

Question 9: Let $A = (a_{ij})$ be a positive semidefinite matrix of real entries. Does the following inequality hold?

$$\text{per} \left((a_{ij}^2) \right) \leq (\text{per } A)^2$$

7 A few more open conjectures on permanent

Below are a few conjectures and research problems that virtually have no progress made in the past years. They are easily stated and understood, but appear to be forgotten. We bring them up again and hope to get renewed attention.

Marcus per-in-per conjecture 1965 [32] *Let A be a positive semidefinite $mn \times mn$ matrix partitioned as $A = (A_{ij})$ in which each A_{ij} is $n \times n$. Let P be the $m \times m$ matrix whose (i, j) entry is $\text{per } A_{ij}$. Then*

$$\text{per } A \geq \text{per } P$$

If A is positive definite, then equality occurs if and only if all $A_{ij} = 0, i \neq j$.

The case of $m = 2$ is proved by Lieb 1966 [28] and Djoković 1969 [11]. A weaker inequality that $\text{per } A \geq (\text{per } P)/n!$ is shown by Pate 1981 [43]. Pate 1982 [44] also shows that for each m , Marcus per-in-per conjecture is true for n sufficiently large. This may be compared with Thompson's det-in-det theorem 1961 [55] which asserts that the determinant of A is dominated by the determinant of D whose (i, j) entry is $\det A_{ij}$, that is, $\det A \leq \det D$.

Marcus-Minc max-per- U problem 1965 [32] *Let A be an $n \times n$ positive semidefinite matrix. Find the maximum value of $\text{per}(U^*AU)$ for all $n \times n$ unitary matrices U .*

Permanent is a continuous function on the compact set of unitary matrices. The minimum and maximum values are achievable. Some upper bounds in terms of eigenvalues and trace of the matrix A have been obtained. For instance,

$$\text{per}(U^*AU) \leq \frac{1}{n} \sum_{i=1}^n \lambda_i^n$$

where λ_i are the eigenvalues of the $n \times n$ positive semidefinite matrix A (see, e.g., [31]). It was conjectured that the maximum of $\text{per}(U^*AU)$ is attained when all the main diagonal entries of U^*AU are all equal (see [41, p. 132]). However, this is false; see a counterexample in [13] or [9, p. 331]. The problem is studied with partial solutions by Drew and Johnson 1989 [12] and Grone et al 1986 [19].

Foregger per- k conjecture 1978 [39, p. 157]. *Let n be a given positive integer. Then there exists a positive integer k (solely depending on n) such that for all $n \times n$ doubly stochastic matrices A*

$$\text{per}(A^k) \leq \text{per } A$$

Chang 1984 [7] and 1990 [8] confirmed the conjecture under certain conditions. For instance, if $\frac{1}{2} < \text{per } A < 1$, then $\text{per}(A^m) < \text{per } A$ for any integer $m \geq 2$. No other progress is seen.

Bapat-Sunder per-max conjecture 1986 [4, Conjecture 3]. *Let A be an $n \times n$ positive semidefinite matrix. Denote by $A(i, j)$ the $(n-1) \times (n-1)$ submatrix of A obtained by deleting the i th row and j th column of A . Then $\text{per } A$ is the maximum eigenvalue of the matrix $(a_{ij} \text{per } A(i, j))$.*

This statement is weaker than the POT conjecture which is true for $n \leq 3$. Hence, the Bapat-Sunder per-max conjecture holds true when the size of the matrix is no more than 3. Other than this, nothing else is known.

For nonnegative $A \in M_n$ and $B \in M_m$, Brualdi 1966 [6] showed that

$$(\text{per } A)^m (\text{per } B)^n \leq \text{per}(A \otimes B)$$

Does this also hold for positive semidefinite matrices?

Liang-So-Zhang per-tensor product conjecture 1992 [59]. *Let A and B be $n \times n$ and $m \times m$ positive semidefinite matrices, respectively. Then*

$$(\text{per } A)^m (\text{per } B)^n \leq \text{per}(A \otimes B)$$

By Brualdi’s result, the conjecture holds true for positive semidefinite matrices of nonnegative entries. In particular, if A and B are square matrices of 1s, it reduces to the known inequality $(n!)^m (m!)^n \leq (nm)!$.

Liang-So-Zhang per-tensor product conjecture, if proven to be true, would imply a result of Ando 1981 [1]:

$$\max\{(n!)^{-m}, (m!)^{-n}\} (\text{per } A)^m (\text{per } B)^n \leq \text{per}(A \otimes B)$$

which is stronger than a result of Marcus 1966 [29]:

$$(n!)^{-m} (m!)^{-n} (\text{per } A)^m (\text{per } B)^n \leq \text{per}(A \otimes B)$$

A special case of Liang-So-Zhang per-tensor product conjecture is the conjecture of Pate 1984 [45]: If $M \in M_m$ is positive semidefinite and J_k is the all one matrix of size k , then

$$(k!)^m (\text{per } M)^k \leq \text{per}(J_k \otimes M) \tag{19}$$

which has been confirmed for $k = 2$ by Pate 1984 [45] and for $m = 2$ [59].

$$2^m (\text{per } M)^2 \leq \text{per} \begin{pmatrix} M & M \\ M & M \end{pmatrix}$$

It would be interesting for one to consider the case where M is 3×3 in (19) and the case where J_k is 3×3 in (19).

In light of the Chollet conjecture $\text{per}(A \circ B) \leq \text{per } A \text{ per } B$, we have

Liang-So-Zhang Hadamard-Kronecker product conjecture 1992 [59]. *Let A and B be $n \times n$ positive semidefinite matrices. Then*

$$(\text{per}(A \circ B))^n \leq \text{per}(A \otimes B)$$

The above conjecture is proved [59, Theorem 2] for any positive semidefinite $A \in M_n$ and special $B = (b_{ij}) \in M_n$ with all diagonal entries equal (to a , say) and all the entries above the main diagonal equal (to $b_{ij} = b$, say, for all $i < j$).

Drury permanent conjecture 2016 [16]. *Let A be an $n \times n$ positive semidefinite matrix and B be the submatrix of A obtained by deleting the first row and column of A . Let B_{jk} be the submatrix of B obtained by deleting the j th row and k th column of B . Then*

$$(a_{11} \text{per } B)^2 + \left(\sum_{k=2}^n |a_{1k}|^2 \text{per}(B_{kk}) \right)^2 \leq (\text{per } A)^2 \tag{20}$$

Two immediate consequences of (20) are $a_{11} \text{per } B \leq \text{per } A$ (well-known) and

$$\sum_{k=2}^n |a_{1k}|^2 \text{per}(B_{kk}) \leq \text{per } A \tag{21}$$

(21) itself is an interesting open problem. Chollet conjecture follows from (20).

Acknowledgement: I am grateful to R. Bapat, R. Brualdi, G.-S. Cheon, S. Drury, and V. Shchesnovich for valuable comments. I am especially indebted to Prof. Drury for his help with the counterexamples. The project was partially supported by National Natural Science Foundation of China (No. 11571220) through Shanghai University.

References

- [1] T. Ando, Inequalities for permanents, *Hokkaido Math. J.* 10 (1981), Special Issue, 18–36.
- [2] R.B. Bapat, Recent developments and open problems in the theory of permanents, *Math. Student* 76 (2007), no. 1-4, 55–69.
- [3] R.B. Bapat and V.S. Sunder, On majorization and Schur products, *Linear Algebra Appl.* 72 (1985) 107–117.
- [4] R.B. Bapat and V.S. Sunder, An extremal property of the permanent and the determinant, *Linear Algebra Appl.* 76 (1986) 153–163.
- [5] L.B. Beasley, An inequality on permanents of Hadamard products, *Bull. Korean Math. Soc.* 37 (2000), no. 3, 633–639.
- [6] R. Brualdi, Permanent of the direct product of matrices, *Pacific J. Math.* 16 (1966) 471–482.
- [7] D.K. Chang, A note on a conjecture of T.H. Foregger, *Linear Multilinear Algebra* 15 (1984), no. 3-4, 341–344.
- [8] D.K. Chang, On two permanental conjectures, *Linear Multilinear Algebra* 26 (1990), no. 3, 207–213.
- [9] G.-S. Cheon and I.-M. Wanless, An update on Minc’s survey of open problems involving permanents, *Linear Algebra Appl.* 403 (2005) 314–342.
- [10] J. Chollet, Unsolved Problems: Is There a Permanental Analogue to Oppenheim’s Inequality? *Amer. Math. Monthly* 89 (1982), no. 1, 57–58.
- [11] D.Z. Djoković, Simple proofs of a theorem on permanents, *Glasgow Math. J.* 10 (1969) 52–54.
- [12] J. Drew and C. Johnson, The maximum permanent of a 3-by-3 positive semidefinite matrix, given the eigenvalues, *Linear Multilinear Algebra* 25 (1989), no. 3, 243–251.
- [13] J. Drew and C. Johnson, Counterexample to a conjecture of Mehta regarding permanental maximization, *Linear Multilinear Algebra* 25 (1989), no. 3, 253–254.
- [14] S. Drury, A counterexample to a question of Bapat and Sunder, *Electronic Journal of Linear Algebra*, Volume 31, pp. 69-70 (2016).
- [15] S. Drury, A real counterexample to two inequalities involving permanents, *Mathematical Inequalities & Applications*, in press.
- [16] S. Drury, Two open problems on permanents, a private communication, May 30, 2016.
- [17] R.J. Gregorac and I.R. Hentzel, A note on the analogue of Oppenheim’s inequality for permanents, *Linear Algebra Appl.* 94 (1987) 109–112.
- [18] R. Grone, An inequality for the second immanant, *Linear Multilinear Algebra* 18 (1985), no. 2, 147–152.
- [19] R. Grone, C. Johnson, E. de S? and H. Wolkowicz, A note on maximizing the permanent of a positive definite Hermitian matrix, given the eigenvalues, *Linear Multilinear Algebra* 19 (1986), no. 4, 389–393.
- [20] R. Grone and R. Merris, Conjectures on permanents, *Linear Multilinear Algebra* 21 (1987), no. 4, 419–427.
- [21] R. Grone, S. Pierce and W. Watkins, Extremal correlation matrices, *Linear Algebra Appl.* 134 (1990) 63–70.
- [22] P. Heyfron, Immanant dominance orderings for hook partitions, *Linear Multilinear Algebra* 24 (1988), no. 1, 65–78.
- [23] R.A. Horn, The Hadamard product, *Proceedings of Symposia in Applied Mathematics*, Vol. 40, edited by C.R. Johnson, Amer. Math. Soc., Providence, RI, pp. 87–169, 1990.
- [24] G. James, Immanants, *Linear Multilinear Algebra* 32 (1992), no. 3-4, 197–210.
- [25] G. James and M. Liebeck, Permanents and immanants of Hermitian matrices, *Proc. London Math. Soc.* (3) 55 (1987), no. 2, 243–265.
- [26] C.R. Johnson, The permanent-on-top conjecture: a status report, in *Current Trends in Matrix Theory*, edited by F. Uhlig and R. Grone, Elsevier Science Publishing Co., New York, pp. 167–174, 1987.
- [27] C.R. Johnson and F. Zhang, Erratum: The Robertson-Taussky Inequality Revisited, *Linear Multilinear Algebra* 38 (1995) 281–282.
- [28] E.H. Lieb, Proofs of some conjectures on permanents, *J. Math. and Mech.* 16 (1966) 127–134.
- [29] M. Marcus, Permanents of direct products, *Proc. Amer. Math. Soc.* 17 (1966) 226–231.
- [30] M. Marcus, *Finite Dimensional Multilinear Algebra. Part I. (Pure and applied mathematics 23)*, Marcel Dekker, New York, 1973.
- [31] M. Marcus and H. Minc, Inequalities for general matrix functions, *Bull. Amer. Math. Soc.* 70 (1964) 308–313.
- [32] M. Marcus and H. Minc, Permanents, *Amer. Math. Monthly* 72 (1965) 577–591.
- [33] M. Marcus and M. Sandy, Bessel’s inequality in tensor space, *Linear Multilinear Algebra* 23 (1988), no. 3, 233–249.
- [34] R. Merris, Extensions of the Hadamard determinant theorem, *Israel J. Math.* 46 (1983), no. 4, 301–304.
- [35] R. Merris, The permanental dominance conjecture, *Current trends in matrix theory : proceedings of the Third Auburn Matrix Theory Conference, March 19-22, 1986 at Auburn University, Auburn, Alabama, U.S.A.*, editors, Frank Uhlig, Robert Grone.

- [36] R. Merris, The permanent dominance conjecture, in *Current Trends in Matrix Theory*, edited by F. Uhlig and R. Grone, Elsevier Science Publishing Co., New York, pp. 213–223, 1987.
- [37] R. Merris, *Multilinear Algebra*, Gordon & Breach, Amsterdam, 1997.
- [38] R. Merris and W. Watkins, Inequalities and identities for generalized matrix functions, *Linear Algebra Appl.* 64 (1985) 223–242.
- [39] H. Minc, *Permanents*, Addison-Wesley, New York, 1978.
- [40] H. Minc, Theory of permanents 1978-1981, *Linear Multilinear Algebra* 12 (1983) 227–263.
- [41] H. Minc, Theory of permanents 1982-1985, *Linear Multilinear Algebra* 21 (1987) 109–148.
- [42] A. Oppenheim, Inequalities Connected with Definite Hermitian Forms, *J. London Math. Soc.* S1-5, no. 2, 114–119.
- [43] T.H. Pate, An extension of an inequality involving symmetric products with an application to permanents, *Linear Multilinear Algebra* 10 (1981), no. 2, 103–105.
- [44] T.H. Pate, Inequalities relating groups of diagonal products in a Gram matrix, *Linear Multilinear Algebra* 11 (1982), no. 1, 1–17.
- [45] T.H. Pate, An inequality involving permanents of certain direct products, *Linear Algebra Appl.* 57 (1984) 147–155.
- [46] T.H. Pate, Permanent dominance and the Soules conjecture for certain right ideals in the group algebra, *Linear Multilinear Algebra* 24 (1989), no. 2, 135–149.
- [47] T.H. Pate, Partitions, irreducible characters, and inequalities for generalized matrix functions, *Trans. Amer. Math. Soc.* 325 (1991), no. 2, 875–894.
- [48] T.H. Pate, Row appending maps, ζ -functions, and immanant inequalities for Hermitian positive semi-definite matrices, *Proc. London Math. Soc.* (3) 76 (1998), no. 2, 307–358.
- [49] T.H. Pate, Tensor inequalities, ζ -functions and inequalities involving immanants, *Linear Algebra Appl.* 295 (1999), no. 1-3, 31–59.
- [50] S. Pierce, Permanents of correlation matrices, in *Current Trends in Matrix Theory*, edited by F. Uhlig and R. Grone, Elsevier Science Publishing Co., New York, pp. 247–249, 1987.
- [51] I. Schur, Über endliche Gruppen und Hermitesche Formen (German), *Math. Z.* 1 (1918), no. 2-3, 184–207.
- [52] V.S. Shchesnovich, The permanent-on-top conjecture is false, *Linear Algebra Appl.* 490 (2016) 196–201.
- [53] G. Soules, *Matrix functions and the Laplace expansion theorem*, Ph.D. Dissertation, University of California - Santa Barbara, July, 1966.
- [54] G. Soules, An approach to the permanent-dominance conjecture, *Linear Algebra Appl.* 201 (1994) 211–229.
- [55] R.C. Thompson, A determinantal inequality, *Canad. Math. Bull.*, vol.4, no.1, Jan. 1961, pp. 57–62.
- [56] J.H. van Lint and R.M. Wilson, *A Course in Combinatorics*, Cambridge University Press, 2001.
- [57] F. Zhang, Notes on Hadamard products of matrices, *Linear Multilinear Algebra* 25 (1989) 237–242.
- [58] F. Zhang, An analytic approach to a permanent conjecture, *Linear Algebra Appl.* 438 (2013) 1570–1579.
- [59] F. Zhang, J. Liang, and W. So, Two conjectures on permanents in *Report on Second Conference of the International Linear Algebra Society (Lisbon, 1992)* by J.A.Dias da Silva, *Linear Algebra Appl.* 197/198 (1994) 791–844.